

AS



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number: **0 533 242 A2**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **92202722.2**

(22) Date of filing: **08.09.92**

(51) Int. Cl.⁵: **C07D 295/22, C07D 207/26, C07D 233/54, C07D 453/02, C07D 303/34, C07D 207/16, C07D 211/60, C07D 211/62, A61K 31/495, C07D 213/38**

(30) Priority: **16.09.91 US 760422**

(43) Date of publication of application:
24.03.93 Bulletin 93/12

(64) Designated Contracting States:
CH DE FR GB IT LI NL


(71) Applicant: **MERCK & CO. INC.**
126, East Lincoln Avenue P.O. Box 2000
Rahway New Jersey 07065-0900(US)

(72) Inventor: **Bock, Mark G.**
1603 Leon Drive
Hatfield, PA 19440(US)
Inventor: **Williams, Peter D.**
260 Shady Nook Road
Harleysville, PA 19438(US)

(74) Representative: **Thompson, John Dr. et al**
Merck & Co., Inc. European Patent
Department Terlings Park Eastwick Road
Harlow, Essex CM20 2QR (GB)

(54) **Substituted derivatives of piperazinylcamphorsulfonyl oxytocin antagonists.**

(57) **Compounds of the formula:**



wherein R is oxime, hydroxy, halogen, cyclic epoxide, alkylcarbonyloxy, alkoxy carbonylalkoxy, trihaloalkylsulfonyloxy, or unsubstituted or substituted oxo, where said substituent is unsubstituted or substituted saturated 6-membered heterocyclic rings having 1 or 2 heteroatoms which are N, and said substituent is alkyl.

The compounds of formula I are oxytocin antagonists useful in the treatment of preterm labor, dysmenorrhea and for the stoppage of labor preparatory to cesarean delivery. Also disclosed are pharmaceutical compositions containing these compounds, methods of their use and methods of their preparation.

EP 0 533 242 A2

FIELD OF THE INVENTION

The present invention provides novel compounds, novel compositions, methods of their use and methods of their manufacture, such compounds generally pharmacologically useful as agents in obstetric and gynecologic therapy. The aforementioned pharmacologic activities are useful in the treatment of mammals. More specifically, the compounds of the present invention can be used in the treatment of preterm labor, stopping labor preparatory to Cesarean delivery, and in the treatment of dysmenorrhea. At the present time, there is a need in the area of obstetric and gynecologic therapy for such agents.

BACKGROUND OF THE INVENTION

In the field of obstetrics, one of the most important problems is the management of preterm labor. A significant number of the pregnancies progressing past 20 weeks of gestation experience premature labor and delivery, which is a leading cause of neonatal morbidity and mortality. Despite major advances in neonatal care, retention of the fetus in utero is preferred in most instances.

Tocolytic (uterine-relaxing) agents that are currently in use include β_2 -adrenergic agonists, magnesium sulfate and ethanol. Ritodrine, the leading β_2 -adrenergic agonist, causes a number of cardiovascular and metabolic side effects in the mother, including tachycardia, increased renin secretion, hyperglycemia (and reactive hypoglycemia in the infant). Other β_2 -adrenergic agonists, including terbutaline and albuterol have side effects similar to those of ritodrine. Magnesium sulfate at plasma concentrations above the therapeutic range of 4 to 8 mg/dL can cause inhibition of cardiac conduction and neuromuscular transmission, respiratory depression and cardiac arrest, thus making this agent unsuitable when renal function is impaired. Ethanol is as effective as ritodrine in preventing premature labor, but it does not produce a corresponding reduction in the incidence of fetal respiratory distress that administration of ritodrine does.


It has been proposed that a selective oxytocin antagonist would be the ideal tocolytic agent. In the last few years, evidence has accumulated to strongly suggest that the hormone oxytocin is the physiological initiator of labor in several mammalian species including humans. Oxytocin is believed to exert this effect in part by directly contracting the uterine myometrium and in part by enhancing the synthesis and release of contractile prostaglandins from the uterine endometrium/decidua. These prostaglandins may, in addition, be important in the cervical ripening process. By these mechanisms, the process of labor (term and preterm) is initiated by a heightened sensitivity of the uterus to oxytocin, resulting in part as a result of a well-documented increase in the number of oxytocin receptors in this tissue. This "up-regulation" of oxytocin receptors and enhanced uterine sensitivity appears to be due to trophic effects of rising plasma levels of estrogen towards term. By blocking oxytocin, one would block both the direct (contractile) and indirect (enhanced prostaglandin synthesis) effects of oxytocin on the uterus. A selective oxytocin blocker, or antagonist, would likely be more efficacious for treating preterm labor than current regimens. In addition, since oxytocin at term has major effects only on the uterus, such an oxytocin antagonizing compound would be expected to have few, if any, side effects.

The compounds of the present invention can also be useful in the treatment of dysmenorrhea. This condition is characterized by cyclic pain associated with menses during ovulatory cycles. The pain is thought to result from uterine contractions and ischemia, probably mediated by the effect of prostaglandins produced in the secretory endometrium. By blocking both the direct and indirect effects of oxytocin on the uterus, a selective oxytocin antagonist can be more efficacious for treating dysmenorrhea than current regimens.

It is, therefore, a purpose of this invention to provide substances which more effectively antagonize the function of oxytocin in disease states in animals, preferably mammals, especially in humans. It is another purpose of this invention to prepare novel compounds which more selectively inhibit oxytocin. It is still another purpose of this invention to provide a method of antagonizing the functions of oxytocin in disease states in mammals. It is also a purpose of this invention to develop a method of preventing or treating oxytocin-related disorders of preterm labor and dysmenorrhea by antagonizing oxytocin.

It has now been found that compounds of formula I are antagonists of oxytocin and bind to the oxytocin receptor. When the oxytocin receptor is bound by the compounds of the present invention, oxytocin is antagonized by being blocked from its receptor and thus being unable to exert its biologic or pharmacologic effects. These compounds are useful in the treatment and prevention of oxytocin-related disorders of animals, preferably mammals and especially humans. These disorders are primarily preterm labor and dysmenorrhea. The compounds would also find usefulness for stoppage of labor preparation to Cesarean delivery.

The compounds of the present invention are those of the general structural formula:



and the pharmaceutically acceptable salts thereof,

25 X is

- (1) carbonyl or
- (2) sulfonyl;

Y is

- (1) hydrogen,
- 30 (2) alkyl or
- (3) NH;

Z is

- (1) C or
- (2) N;

35 R¹ and R² are independently

- (1) hydrogen,
- (2) halogen,
- (3) alkoxy,
- (4) alkylsulfonyl or
- 40 (5) unsubstituted or substituted alkyl wherein said substituent is
- amino,
- alkyl, or
- dialkylamino;

R³ and R⁴ are independently

- 45 (1) hydrogen,
- (2) alkyl,
- (3) substituted alkyl when said substituent is
- amino
- alkylsulfonyl,
- 50 arylsulfonyl,
- alkylamino, or
- dialkylamino;
- (4) phenylalkyl or
- (5) oxo;

55 R⁵ is

- (1) hydrogen or
- (2) oxo


R⁶ and R⁷ are independently

- (1) hydrogen,
 (2) alkyl or
 (3) joined to form unsubstituted or substituted cycloalkyl where said substituent is hydroxy or hydroxyalkyl;
 5 R^8 , R^9 and R^{10} are independently
 (1) hydrogen,
 (2) oxime,
 (3) hydroxy,
 10 (4) halogen,
 (5) cyclic epoxide,
 (6) alkylcarbonyloxy,
 (7) alkoxycarbonylalkoxy,
 (8) trihaloalkylsulfonyloxy,
 15 (9) unsubstituted or substituted 5 or 6-membered heterocyclic rings containing 1,2 or 3 heteroatoms where said heteroatom is N,S or O and said substituents are one or more of oxo or,
 C_{1-6} alkyl, optionally substituted by
 alkylthio
 20 alkylsulfinyl
 alkylsulfonyl
 cyano
 carboxy,
 hydroxy
 25 amine,
 alkylamine
 dialkylamine
 alkoxy
 aminocarbonyl,
 30 tetrazolyl
 imidazole and
 spiro-piperidinyl where the piperidine N is unsubstituted or substituted by C_{1-6} alkyl;
 (10) unsubstituted, mono- or di-substituted phenyl where said substituents are independently one or more of
 35 halogen,
 carboxy,
 alkoxycarbonyl,
 carboxyalkyl,
 amino,
 40 alkylamino,
 dialkylamino,
 cyano,
 alkylsulfonylamino,
 aminoalkyl,
 45 alkylaminoalkyl,
 dialkylaminoalkyl; and
 m, n, p and q are integers of from 0 to 2.

More particularly preferred compounds are those of the general structural Formula

50

55



and the pharmaceutically acceptable salts thereof.

X is

- (1) carbonyl or
- (2) sulfonyl;

Y is

- (1) hydrogen,
- (2) alkyl or
- (3) NH;

Z is

- (1) C or
- (2) N;

R¹ and R² are independently

- (1) hydrogen,
- (2) halogen,
- (3) alkoxy,
- (4) alkylsulfonyl or
- (5) unsubstituted or substituted alkyl wherein said substituent is amino, alkylamino or dialkylamino;

R³ and R⁴ are independently

- (1) hydrogen,
- (2) alkyl,
- (3) substituted alkyl where said substituent is amino, alkylsulfonyl, arylsulfonyl, alkylamino, or dialkylamino;
- (4) phenylalkyl or
- (5) oxo;

R⁵ is

- (1) hydrogen or
- (2) oxo;

R⁶ and R⁷ are independently

- (1) hydrogen,
- (2) alkyl or

(3) joined to form unsubstituted or substituted cycloalkyl where said substituent is hydroxy or hydroxyalkyl;

R^8 , R^9 and R^{10} are independently

- 5 (1) hydrogen,
- (2) oxime,
- (3) hydroxy,
- (4) halogen,
- (5) oxo,
- 10 (6) cyclic epoxide,
- (7) alkylcarbonyloxy,
- (8) alkoxy carbonylalkoxy,
- (9) trihaloalkylsulfonyloxy,
- (10) unsubstituted or substituted 5-membered heterocyclic rings containing 1 or 2 heteroatoms where
- 15 said heteroatom is N or O and said substituents are oxo or
- C_{1-6} alkyl, optionally substituted by
- carboxy,
- amine,
- 20 aminocarbonyl or
- imidazole; and

m, n, p and q are integers from 0 to 2.


Most preferred are those compounds of the general formula

25

30

35

40



45

and the pharmaceutically acceptable salts thereof, wherein

R' is an optional substituent that, when present, is H or OH;

R is

- 50 (1) hydrogen,
- (2) oxime,
- (3) hydroxy,
- (4) halogen,
- (5) cyclic epoxide,
- (6) alkylcarbonyloxy,
- 55 (7) alkoxy carbonylalkoxy,
- (8) trihaloalkylsulfonyloxy,
- (9) unsubstituted or substituted 5-membered heterocyclic rings containing 1 or 2 heteroatoms where said
- heteroatom is N or O and said substituents are

oxo or

C₁₋₆ alkyl, optionally substituted by

carboxy,

amine,

aminocarbonyl or

imidazole.

Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention which are generally prepared by reacting the free base with a suitable organic or inorganic acid. Representative salts include the following salts:

Acetate	Lactobionate
Benzenesulfonate	Laurate
Benzoate	Malate
Bicarbonate	Maleate
Bisulfate	Mandelate
Bitartrate	Mesylate
Borate	Methylbromide
Bromide	Methylnitrate
Calcium Edetate	Methylsulfate
Camsylate	Mucate
Carbonate	Napsylate
Chloride	Nitrate
Clavulanate	N-methylglucamine ammonium salt
Citrate	
Dihydrochloride	Oleate
Edetate	Oxalate
Edisylate	Pamoate (Embonate)
Estolate	Palmitate
Esylate	Pantothenate
Fumarate	Phosphate/diphosphate
Gluceptate	Polygalacturonate
Gluconate	Salicylate
Glutamate	Stearate
Glycolylarsanilate	Subacetate
Hexylresorcinate	Succinate
Hydrabamine	Tannate
Hydrobromide	Tartrate
Hydrochloride	Teoclate
Hydroxynaphthoate	Tosylate
Iodide	Triethiodide
Isethionate	Valerate
Lactate	

The term "pharmacologically effective amount" shall mean that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by a researcher or clinician.

The term "alkyl" shall mean straight or branched chain alkanes, alkenes and alkynes with one or more degrees of unsaturation at any position on the chain, of one to ten total carbon atoms, or any number within this range.

The term "aryl" shall mean phenyl, naphthyl and fluorenyl.

The term "cycloalkyl" shall mean cyclic rings of alkanes, alkenes or alkynes with one or more degrees of unsaturation at any position of the ring, of three to eight total carbon atoms.

Whenever the terms "alkyl" or "aryl" or either of their prefix roots appear in a name of a substituent (e.g. aralkoxyaryloxy) they shall be interpreted as including those limitations given above for "alkyl" and "aryl". Designated numbers of carbon atoms (e.g. C₁₋₁₀) shall refer independently to the number of carbon atoms in an alkyl or cyclic alkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears

as its prefix root.

Where multiple substituent moieties are disclosed or claimed, the substituted compound can be independently substituted by one or more of the disclosed or claimed substituent moieties, singly or plurally.

5 The term "oxo" shall refer to the substituent =O.

The term "halogen" shall include iodine, bromine, chlorine and fluorine.

The term "preterm labor" shall mean expulsion from the uterus of a viable infant before the normal end of gestation, or more particularly, onset of labor with effacement and dilation of the cervix before the 37th week of gestation. It may or may not be associated with vaginal bleeding or rupture of the membranes.

10 The term "dysmenorrhea" shall mean painful menstruation.

The term "cesarean delivery" shall mean incision through the abdominal and uterine walls for delivery of a fetus.

The term "substituted" shall be deemed to include multiple degrees of substitution by a named substituent.

15 The ability of the compounds of formula I to antagonize oxytocin makes these compounds useful as pharmacologic agents for mammals, especially for humans, for the treatment and prevention of disorders wherein oxytocin may be involved. Examples of such disorders include preterm labor and especially dysmenorrhea. These compounds may also find usefulness for stoppage of labor preparatory to Cesarean delivery.

20 Because of the known relationship of vasopressin to oxytocin, the compounds of the present invention are also useful as vasopressin antagonists. Vasopressin antagonists are useful in the treatment or prevention of disease states involving vasopressin disorders, including their use as diuretics and their use in congestive heart failure.

The compounds of the present invention can be administered in such oral dosage forms as tablets, capsules (each including timed release and sustained release formulations), pills, powders, granules, elixers, tinctures, suspensions, syrups and emulsions. Likewise, they may also be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the compound desired can be employed as a tocolytic agent.

30 The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

35 Oral dosages of the present invention, when used for the indicated effects, will range between about 0.3-6.0 gm/day orally. Intravenously, the most preferred doses will range from 0.1 to about 10 mg/minute during a constant rate infusion. Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, preferred compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

45 In the methods of the present invention, the compounds herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

50 For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

5 Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

10 The compounds of formula I can be prepared readily according to the following reaction Schemes (in which all variables are as defined before) and Examples or modifications thereof using readily available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of variants which are themselves known to those of ordinary skill in this art, but are not mentioned in greater detail.

15 The most preferred compounds of the invention are any or all of those specifically set forth in these examples. These compounds are not, however, to be construed as forming the only genus that is considered as the invention, and any combination of the compounds or their moieties may itself form a genus. The following examples further illustrate details for the preparation of the compounds of the present invention. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds. All temperatures are degrees Celsius unless noted otherwise.

30





35

40




45

50


55

SCHEME 1


II

SCHEME 2**SCHEME 3****SCHEME 4**


SCHEME 5




SCHEME 6



SCHEME 7



SCHEME 8



Abbreviations used in the Examples are as follows:

TEA = triethylamine

DIEA = diisopropylethylamine

BOP = benzotriazolyloxytris(dimethylamino) phosphonium hexafluorophosphate

THF = tetrahydrofuran

DMF = dimethylformamide

LAH = lithium aluminum hydride

TFA = trifluoroacetic acid

HPLC Method A = 15 min. linear gradient

95:5 A:B to 0:100 A:B

A = H₂O containing 0.1% by vol. TFA

B = CH₃CN containing 0.1% by vol. TFA

2.0 mL/min flow rate


12 cm C₁₈ reverse phase column

UV detection (215 nm)

TLC was performed on 20 cm plates coated with silica gel (250 microns) from Analtech.

Example 1

1-((7,7-DIMETHYL-2-OXO-BICYCLO(2.2.1)HEPTAN-1-YL) METHANESULFONYL)-4-(2-METHYLPHENYL)-PIPERAZINE



To a stirred, 0°C solution of 1-(o-tolyl)piperazine hydrochloride (50.0 g; 235 mmol) and TEA (83 mL; 590 mmol) in chloroform (1000 mL) was added (+)-10-camphorsulfonyl chloride (65.5 g; 260 mmol). The solution was stirred at 0°C for 1 h and then at ambient temperature for 3 h. The solution was extracted with 5% aqueous HCl (2 x 500 mL), water (500 mL), and saturated aqueous NaHCO₃ (2 x 500 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The resulting solid was recrystallized from methanol to give the title compound, mp 112-114°C (69 g; 75%).

anal: (C ₂₁ H ₃₀ N ₂ O ₃ S)			
calc.	C, 64.57;	H, 7.74;	N, 7.17
found	C, 64.52;	H, 7.68;	N, 6.99

TLC: R_f 0.49 (75:25 hexane/ethyl acetate)


HPLC (method A): retention time 10.33 min

FAB MS: m/z 391 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.2 (m, 2H), 7.0 (m, 2H), 3.45 (m, 4H), 3.40 (d, J = 16 Hz, 1H), 3.0 (m, 4H), 2.57 (m, 1H), 2.40 (dt, J_d = 14 Hz, J_t = 3 Hz, 1H), 2.30 (s, 3H), 2.10 (m, 2H), 1.96 (d, J = 14 Hz, 1H), 1.67 (m, 1H), 1.44 (m, 1H), 1.18 (s, 3H), 0.91 (s, 3H)

Example 2

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-(1-CYANO)ETHYL-BICYCLO(2.2.1)HEPTAN-1-YL)-METHANESULFONYL)-4-(2-METHYL PHENYL)PIPERAZINE:



To a stirred, -78 °C solution of diisopropylamine (21.0 mL; 150 mmol) in THF (350 mL) was added n-butyllithium (60 mL of a 2.5 M solution in hexane; 150 mmol). The solution was warmed to 0 °C for 15 min, then cooled to -78 °C. A solution of propionitrile (10.1 mL; 141 mmol) in THF (75 mL) was added dropwise, and the resulting solution was stirred at -78 °C for 45 min. A -78 °C solution of 1-((7,7-dimethyl-2-oxo-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (50.0 g; 128 mmol) in THF (350 mL) was added via cannula, and the resulting solution was stirred at -78 °C for 5 min. A solution of 5:1 THF/water (100 mL) was added and the mixture was warmed to ambient temperature. The mixture was diluted with EtOAc (500 mL) and washed with 5% aqueous citric acid (2 x 500 mL), and brine (250 mL). The organic phase was dried (MgSO₄), filtered, and the solvents were removed under reduced pressure to give a foam. The major isomer by TLC was obtained by crystallization from ether, mp 163-165 °C.

anal: (C ₂₄ R ₃₅ N ₃ O ₃ S)			
calc.	C, 64.69;	H, 7.92;	N, 9.43
found	C, 64.72;	H, 7.99;	N, 9.35

TLC: R_f 0.31 (75:25 hexane/ethyl acetate)


HPLC (method A): retention time 10.20 min

FAB MS: m/z 446 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.19 (m, 2H), 3.70 (d, J = 15 Hz, 1H), 3.68 (s, 1H), 3.49 (m, 4 H), 3.38 (d, J = 15 Hz, H), 2.75 (q, J = 7 Hz, 1H), 2.30 (s, 2H), 2.05 (m, 2H), 1.7-1.9 (m, 3H), 1.47 (d, J = 7 Hz, 3H), 1.41 (d, J = 12 Hz, 1H), 1.40 (s, 3H), 1.15 (s, 3H), 1.04 (m, 1H)

Example 3

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)-METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred, -78°C solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-(1-cyano)ethyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (25.0 g; 56.2 mmol) in THF (350 mL) was added dropwise a 1.0 M solution of LAH in THF (170 mL; 170 mmol). The resulting solution was stirred at -78°C for 1 h, and then warmed to 0°C for 3 h. Ether (300 mL) was added, followed by the slow dropwise addition of 5 M NaOH solution (35 mL). The resulting suspension was warmed to ambient temperature and stirred for 1 h. EtOAc (250 mL) was added and stirring was continued for 30 min. The solids were removed by filtration through Celite and washed with EtOAc. The filtrate solvents were removed under reduced pressure to give a foam. The title compound was obtained by crystallization from methanol, mp $172-174^{\circ}\text{C}$ (17.2 g; 68%).

anal: (C ₂₄ H ₃₃ N ₃ O ₃ S)			
calc.	C, 64.11;	H, 8.74;	N, 9.35
found	C, 64.09;	H, 8.88;	N, 9.31

TLC: R_f 0.50 (95:5:0.5 CHCl₃/MeOH/NH₄OH)


HPLC (method A): retention time 9.80 min

FAB MS: m/z 450 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.20 (m, 2H), 7.05 (m, 2H), 2.32 (s, 3H), 1.13 (d, J = 6 Hz, 3H), 1.11 (s, 3H), 1.02 (s, 3H)

Example 4

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(L-PROLYL)AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)-METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-*exo*-hydroxy-2-*endo*-2-(1-amino)propyl-(2.2.1)bicycloheptan-1-yl)-methanesulfonyl)-4-(2-methylphenyl) piperazine (2.00 g; 4.45 mmol) in DMF (30 mL) was added $N\alpha$ -Fmoc-L-proline (1.58 g; 4.68 mmol), BOP (2.17 g; 4.90 mmol), and DIEA (1.71 mL; 9.80 mmol). After 16 h, diethylamine (6 mL) was added and the solution was stirred at ambient temperature for 3 h. The solvents were removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

anal: (C ₂₉ H ₄₆ N ₄ O ₄ S)			
calc.	C, 52.48;	H, 6.50;	N, 7.56
found	C, 52.46;	H, 6.50;	N, 7.69

1.7 TFA, 0.05 H₂O

TLC: R_f 0.45 (90:10:1 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 8.60 min

FAB MS: m/z 547 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.55 (br t, 1H), 7.18 (m, 2H), 7.03 (m, 2H), 2.31 (s, 3H), 1.14 (s, 3H), 1.02 (s, 3H), 0.99 (d, J = 7 Hz, 3H)

Example 5

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(L-N-(ETHOXYCARBONYLPROPYL)PROLYL)AMINO)-PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL) PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-*exo*-hydroxy-2-*endo*-2-(1-(L-prolyl)amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl) piperazine (1.50 g; MW = 679; 2.21 mmol) in DMF (15 mL) was added ethyl 4-bromobutyrate (538 mg; 2.76 mmol), and DIEA (1.15 mL; 6.63 mmol). After 72 h at ambient temperature, the solvent was removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of the title compound was obtained as a lyophilized powder.

anal: (C ₃₅ H ₅₅ N ₄ O ₆ S)			
calc.	C, 51.99;	H, 6.48;	N, 6.17
found	C, 52.01;	H, 6.33;	N, 6.17

2.1 TFA, 0.1 H₂O

TLC: R_f 0.40 (95:5 CHCl₃:MeOH)


HPLC (method A): retention time 10.23 min

FAB MS: m/z 661 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 8.55 (m, 1H), 7.20 (m, 2H), 7.08 (m, 2H), 2.35 (s, 3H), 1.25 (t, J = 6 Hz, 3H), 1.14 (s, 3H), 1.03 (overlapping s and d, 6H)

Example 6

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(L-N-(3-CARBOXYPROPYL)PROLYL)AMINO) PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(L-N-(ethoxycarbonylpropyl) prolyl)-amino)propyl-(2.2.1)bicycloheptan-1-yl) methanesulfonyl)-4-(2-methylphenyl)piperazine (1.00 g; MW = 909; 1.10 mmol) in THF (15 mL) was added 1 M NaOH solution (1.0 mL; 4.0 mmol) until a pH 10 solution persisted for 1 h. The solution was acidified to pH 7 by addition of citric acid and the solvents were removed under reduced pressure. The residue was dissolved in dichloromethane (75 mL) and washed with water (3 x 25 mL), dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was lyophilized from dioxane-water to give the title compound as a white powder.

anal: (C ₃₃ H ₅₂ N ₄ O ₆ S)			
calc.	C, 59.78;	H, 8.25;	N, 6.94
found	C, 59.86;	H, 7.98;	N, 6.92

0.1 Na citrate, 1.65 dioxane

TLC: R_f (80:20:2 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 9.24 min

FAB MS: m/z 633 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.55 (br s, 1H), 7.18 (m, 2H), 7.03 (m, 2H), 2.31 (s, 3H), 1.15 (s, 3H), 1.04 (s, 3H), 0.98 (d, J = 6 Hz, 3H)

Example 7

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(4(5)-IMIDAZOLYLACETYL)AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2- exo-hydroxy-2-endo-2-(1-amino)propyl-(2.2.1)bicyclo-heptan -1-yl)-methanesulfonyl)-4-(2-methylphenyl)piperazine (1.50 g; 3.34 mmol) in DMF (15 mL) was added 4(5)-

imidazol acetic acid hydrochloride (679 mg; 4.18 mmol), BOP (1.85 g; 4.18 mmol), and DIEA (2.18 mL; 12.5 mmol). After 16 h, the solvent was removed under reduced pressure. The residue was dissolved in EtOAc (100 mL) and washed with saturated aqueous NaHCO₃ solution (2 x 50 mL) and water (2 x 50 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 92:8:0.8 as eluant. The title compound crystallized from EtOAc, mp 159-163°C.

anal: (C ₂₉ H ₄₃ N ₅ O ₄ S)			
calc.	C, 62.45;	H, 7.77;	N, 12.56
found	C, 62.88;	H, 7.68;	N, 12.79

TLC: R_f 0.4 (90:10:1 CHCl₃/MeOH/NH₄OH)


HPLC (method A): retention time 8.72 min

FAB MS: m/z 558 (M⁺ + H)

¹H NMR (CDCl₃): δ 7.57 (s, 1H), 7.2 (m, 3H), 7.0 (m, 2H), 6.88 (s, 1H), 3.55 (m, 2H), 3.4 (m, 5H), 2.95 (m, 4H), 2.87 (d, J = 15 Hz, 1H), 2.31 (s, 3H), 1.71 (t, J = 4 Hz, 1H), 1.52 (d, J = 13 Hz, 1H), 1.15 (s, 3H), 1.03 (s, 3H), 0.97 (d, J = 6 Hz, 3H)

Example 8

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-QUINUCLIDIN-3-YL-CARBONYL)AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-amino)propyl-(2.2.1)bicycloheptan-1-yl)-methanesulfonyl)-4-(2-methylphenyl) piperazine (2.00 g; 4.45 mmol) in DMF (50 mL) was added quinuclidine-3-carboxylic acid hydrochloride (938 mg; 4.90 mmol), BOP (2.17 g; 4.90 mmol), and DIEA (2.56 mL; 14.7 mmol). After 16 h, the solvent was removed under reduced pressure. The residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 1% acetic acid. The acetate salt of the title compound (1:1 mixture of diastereomers) was obtained as a lyophilized powder.

anal: (C ₃₂ H ₅₀ N ₄ O ₄ S)			
calc.	C, 60.39;	H, 8.58;	N, 8.39
found	C, 60.41;	H, 8.19;	N, 8.58

0.8 CH₃CO₂H, 1.85 H₂O

TLC: R_f 0.65 (80:20:2 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 8.68 min

FAB MS: m/z 587 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.19 (m, 2H), 7.02 (m, 2H), 2.30 (s, 3H), 1.16 (s, 3H), 1.03 (overlapping s and d, 6H)

Example 9

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(1-CARBOXYMETHYLQUINUCLIDIN-3-YL-CARBOXYL)-AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)-PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(quinuclidin-3-yl-carbonyl)amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (1.50 g; MW = 668; 2.25 mmol) in DMF (30 mL) was added iodoacetic acid (543 mg; 2.92 mmol) and DIEA (0.43 mL; 2.48 mmol). After 16 h, TLC showed complete consumption of starting material. The solvent was removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 1% acetic acid. The title compound, as a 1:1 mixture of diastereomers, was obtained as a lyophilized powder.

anal: (C ₃₄ H ₅₂ N ₄ O ₄ S)			
calc.	C, 60.52;	H, 8.18;	N, 8.04
found	C, 60.52;	H, 7.98;	N, 8.15

0.55 CH₃CO₂H, 0.95 H₂O

TLC: R_f 0.20 (80:20:2 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 8.73 min

FAB MS: m/z 647 (M⁺ + H)

¹H NMR (TFA salt; 400 MHz, CDCl₃): δ 7.46 (br s, 1H), 7.19 (m, 2H), 7.02 (m, 2H), 2.30 (s, 3H), 1.13 (s, 3H), 1.02 (s, 3H), 0.98 (d, J = 6 Hz, 3H)

Example 10

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(2-METHOXYCARBONYLETHYL)AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-amino)propyl-(2.2.1)bicycloheptan-1-yl)-methanesulfonyl)-4-(2-methylphenyl)piperazine (100 mg; 0.22 mmol) in 1:1 DMF-MeOH (3 mL) was added methyl acrylate (0.020 mL; 0.22 mmol). After 16 h, the solvents were removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of the title compound was obtained as a lyophilized powder.

anal: (C ₂₈ H ₄₅ N ₃ O ₅ S)			
calc.	C, 53.03;	H 6.88;	N, 6.06
found	C, 53.01;	H, 6.90;	N, 6.01

1.3 TFA, 0.5 H₂O

TLC: R_f 0.35 (95:5 CHCl₃:MeOH)


HPLC (method A): retention time 9.04 min

FAB MS: m/z 536 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.20 (m, 2H), 7.03 (m, 2H), 3.72 (s, 3H), 2.32 (s, 3H), 1.19 (d, J=6 Hz, 3H), 1.15 (s, 3H), 0.98 (s, 3H)

EXAMPLE 11

1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-bis-(2-methoxycarbonyl)ethyl)amino)propyl-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-amino)-propyl-(2.2.1)bicycloheptan-1-yl)-methanesulfonyl)-4-(2-methylphenyl) piperazine (100 mg; 0.22 mmol) in 1:1 DMF-MeOH (3 mL) was added methyl acrylate (0.080 mL; 0.89 mmol). After 16 h, the solvents were removed under reduced pressure and the residue was purified by pressurized silica gel chromatography using 3:1 hexane-ethyl acetate as eluant. The title compound was obtained as a foam from hexane.

Anal: (C ₃₂ H ₅₁ N ₃ O ₇ S)			
Calc:	C 61.81,	H 8.27,	N 6.76
Found:	C 61.55,	H 8.13,	N 6.55

TLC: R_f 0.40 (1:3 EtOAc:hexanes)


HPLC (method A): retention time 9.71 min

FAB MS: m/z 622 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.19 (m, 2H), 7.02 (m, 2H), 3.66 (s, 6H), 2.31 (s, 3H), 1.13 (s, 3H), 1.00 (overlapping a and d, 6H)

EXAMPLE 12

1-((7,7-dimethyl-2-exo-hydroxy-2-endo-ethenyl-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine




To a -78°C stirred 1.0 M solution of vinyl magnesium chloride in THF (25 mL; 25 mmol) was added a -78°C solution of 1-((7,7-dimethyl-2-oxo-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (5.00 g; 12.8 mmol) in THF (100 mL) via cannula. The resulting solution was stirred under argon overnight, allowing the cooling bath to warm to ambient temperature. The reaction was quenched by addition of 2% aqueous HCl (50 mL), and the mixture was partitioned between ethyl acetate and water. The organic phase was washed with aqueous NaHCO_3 and brine, dried over MgSO_4 , and filtered. The solvents were removed under reduced pressure and the residue was purified by pressurized silica gel chromatography using 4:1 hexane-ethyl acetate as eluant. The title compound was obtained as a white foam from ether.

Anal: $(\text{C}_{23}\text{H}_{34}\text{N}_2\text{O}_3\text{S}) 0.06 \text{ H}_2\text{O}$			
Calc:	C 65.82,	H 8.19,	N 6.67
Found:	C 65.99,	H 8.42,	N 6.63

TLC: R_f 0.36 (1:5 EtOAc:hexanes)
HPLC (method A): retention time 11.41 min
FAB MS: m/z 419 ($\text{M}^+ + \text{H}$)
 ^1H NMR (400 MHz, CDCl_3): δ 7.20 (m, 2H), 7.02 (m, 2H), 6.48 (dd, 1H), 5.30 (d, 1H), 5.17 (d, 1H), 2.32 (s, 3H), 1.22 (s, 3H), 0.94 (s, 3H).

EXAMPLE 13

1-((7,7-dimethyl-2-(2-chloro)ethylidene-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)-piperazine




To a 0°C stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-ethenyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (2.90 g; 6.94 mmol) in THF (100 mL) was added triethylamine (1.50 mL; 10.7 mmol) and DMF (0.58 mL; 7.5 mmol). Thionyl chloride (0.66 mL; 9.1 mmol) was added dropwise, and the resulting solution was stirred for 18 h, allowing the cooling bath to warm ambient temperature. The solvents were removed under reduced pressure and the residue was dissolved in ethyl acetate (150 mL) and washed with 5% aqueous HCl (75 mL), water (75 mL) and aqueous NaHCO₃ (100 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 4:1 hexane-ethyl acetate as eluant. The title compound was obtained as a white foam.

Anal: (C ₂₃ H ₃₃ ClN ₂ O ₂ S) 0.6 H ₂ O			
Calc:	C 65.82,	H 8.19,	N 6.67
Found:	C 65.99,	H 8.42,	N 6.63

¹H NMR (400 MHz, CDCl₃): δ 7.20 (m, 2H), 7.03 (m, 2H), 5.87 (m, 1H), 2.32 (s, 3H), 1.00 (s, 3H), 0.82 (s, 3H)

EXAMPLE 14

1-((7,7-dimethyl-2-(2-isobutylamino)ethylidene-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine



To a stirred solution of 1-((7,7-dimethyl-2-(2-chloro)ethylidene)-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (200 mg; 0.46 mmol) in MeOH (2 mL) was added isobutylamine (0.5 mL; 5 mmol). After being stirred for 18 h, the solvents were removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The

5 TFA salt of the title compound was obtained as a lyophilized powder.

TLC: R_f 0.30 (95:5:0.5 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 9.78 min

FAB MS: m/z 474 ($M^+ + H$)

¹H NMR (400 MHz, CD₃OD): δ 7.20 (m, 3H), 7.03 (t, 1H), 5.78 (m, 1H), 2.35 (s, 3H), 1.13 (d, $J = 7$ Hz, 6H),
10 1.12 (s, 3H), 0.88 (s, 3H)

EXAMPLE 15

1-((7,7-dimethyl-2-(2-azido)ethylidene-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)-
15 piperazine



To a stirred solution of 1-((7,7-dimethyl-2-*exo*-hydroxy-2-(2-chloro)ethylidene-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl) piperazine (3.58 g; 8.19 mmol) in DMSO (50 mL) and THF (45 mL) was added a solution of sodium azide (5.3 g; 82 mmol) in water (20 mL). After 24 h, the solvents were removed under reduced pressure, the residue was suspended in dichloromethane (100 mL) and washed with water (3
35 x 50 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure to give a solid.

Anal: (C ₂₃ H ₃₃ N ₅ O ₂ S)			
Calc:	C 62.27,	H 7.50,	N 15.79
Found:	C 62.41,	H 7.54,	N 15.60

TLC: R_f 0.75 (70:30 hexane-ethylacetate)


HPLC (method A): retention time 12.50 min

45 FAB MS: m/z 444 ($M^+ + H$)

¹H NMR (300 MHz, CDCl₃): δ 7.20 (m, 2H), 7.02 (m, 2H), 5.79 (m, 1H), 2.32 (s, 3H), 0.85 (s, 3H)

EXAMPLE 16

1-((7,7-dimethyl-2-(2-amino)ethylidenebicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-(2-azido)ethylidene-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (3.85 g; 8.69 mmol) in THF (150 mL) and water (3 mL) was added triphenylphosphine (2.50 g; 9.56 mmol). After 14 h, the solvents were removed under reduced pressure. The residue was dissolved in ethyl acetate (150 mL) and extracted with 5% aqueous HCl (3 x 75 mL). The combined acid extracts were washed with ethyl acetate (50 mL) and then made basic by adding solid sodium hydroxide to pH 12. The aqueous phase was extracted with chloroform (3 x 50 mL) and the combined organic phases were dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using a gradient elution of 99:1 to 85:15 chloroform-methanol. The title compound was obtained as a solid.

Anal: (C₂₃H₃₅N₃O₂S) 0.5 H₂O

Calc:	C 64.75;	H 8.51;	N 9.85;
Found:	C 64.59;	H 7.51;	N 9.71


TLC: R_f 0.56 (95:5:0.5 CHCl₃-MeOH-NH₄OH)

HPLC (method A): retention time 10.38 min

FAB MS: m/z 418 (M⁺ + H)

EXAMPLE 17

1-((7,7-dimethyl-2-(2-(4(5)-imidazolylacetyl)amino)ethylidenebicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-(2-amino)ethylidin-(2.2.1)bicycloheptan-1-yl)-methanesulfonyl)-4-(2-methylphenyl)piperazine (0.20 g; 0.48 mmol) in DMF (5 mL) was added BOP (265 mg; 0.60 mmol), 4-imidazoleacetic acid hydrochloride (115 mg; 0.72 mmol) and DIEA (0.38 mL; 2.2 mmol). After 14 h, the solvents were removed under reduced pressure, the residue was suspended in ethyl acetate (50 mL) and washed with aqueous NaHCO₃ (2 x 25 mL) and water (2 x 25 mL). The organic phase was dried (MgSO₄), filtered and the solvent was removed under reduced pressure. The residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of the title compound was obtained as a lyophilized powder.

Anal: (C ₂₈ H ₃₉ N ₅ O ₃ S); 0.5 H ₂ O; 2.0 TFA;			
Calc:	C 50.38;	H 5.55;	N 9.18
Found:	C 50.40;	H 5.55;	N 9.40

TLC: R_f 0.42 (95:5:0.5 CHCl₃-MeOH-NH₄OH)


HPLC (method A): retention time 8.76 min.

FAB MS: m/z 526 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 8.40 (s, 1H), 7.58 (br m, 1H), 7.22 (m, 3H), 7.10 (m, 2H), 5.57 (br t, 1H), 2.37 (s, 3H), 0.97 (s, 3H), 0.76 (s, 3H)

Example 18

1-((7,7-DIMETHYL-2-SPIRO-EPOXY-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred 0 °C suspension of trimethylsulfoxonium iodide (6.78 g; 30.8 mmol) in THF (100 mL) was added n-butyllithium (11.1 mL of a 2.5 M solution in hexane; 27.7 mmol). After 4 h at 0 °C, a solution of 1-((7,7-dimethyl-2-oxo-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (8.00 g; 20.5 mmol) in THF (50 mL). The resulting solution was stirred at 0 °C for 2 h, and then at ambient temperature for 18 h. The solvents were removed under reduced pressure, the residue was dissolved in ethyl acetate (150 mL) and washed with water (2 x 50 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The resulting solid was recrystallized from ether to give the title compound, as white needles.

anal: (C ₂₂ H ₃₂ N ₂ O ₂ S)			
calc.	C, 65.31;	H, 7.97;	N, 6.92
found	C, 65.09;	H, 7.99;	N, 6.86

0.5 H₂O

TLC: R_f 0.62 (4:1 hexane-ethyl acetate)

HPLC (method A): retention time 11.50 min


FAB MS: m/z 405 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.20 (m, 2H), 7.02 (m, 2H), 3.20 (d, J=5.4 Hz, 1H), 2.70 (d, J=5.4 Hz, 1H),

2.30 (s, 3H), 1.00 (s, 3H), 0.99 (s, 3H)

Example 19

5 1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-ISOBUTYLAMINOMETHYL-BICYCLO(2.2.1)HEPTAN-1-YL)-METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-(spiro-epoxy)-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (200 mg; 0.495 mmol) in MeOH (3 mL) was added isobutylamine (0.5 mL; 5 mmol). After being stirred for 18 h, the solvents were removed under reduced pressure and the residue was purified by pressurized silica gel column chromatography using 98:2:0.2 chloroform-methanol-NH₄OH as eluant. The product was dissolved in methanol and to it was added several drops of 5% aqueous HCl. The solvents were removed under reduced pressure and the residue was triturated in ether to give the hydrochloride salt of the title compound as a white powder.

anal: (C ₂₆ H ₄₃ N ₃ O ₃ S)			
calc.	C, 57.00;	H, 8.76;	N, 7.67
found	C, 57.03;	H, 8.84;	N, 7.61

1.0 HCl, 1.8 H₂O

TLC (free base): R_f 0.20 (3:1 hexane-ethyl acetate)

HPLC (method A): retention time 9.54 min

40 FAB MS: m/z 478 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.20 (m, 2H), 7.02 (m, 2H), 2.30 (s, 3H), 1.10 (s, 3H), 0.95 (s, 3H), 0.90 (two doublets, 6H)


45

50

55

Example 20

1-((7,7-DIMETHYL-2-METHOXYCARBONYL-BICYCLO (2.2.1)HEPT-2-EN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:




To a stirred, 0 °C solution of 1-((7,7-dimethyl-2-oxo-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (10.0 g; 25.6 mmol) in dichloromethane (500 mL) was added 2,6-di-*t*-butyl-4-methylpyridine (7.8 g; 38 mmol) and trifluoromethanesulfonic anhydride (5.4 mL; 32 mmol). The cooling bath was removed and the solution was stirred for 18 h. The mixture was filtered and the filtrate was washed with 5% aqueous HCl (2 x 100 mL), water (100 mL), and aqueous NaHCO₃ (2 x 100 mL). The organic phase was dried (MgSO₄), filtered and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 9:1 hexane-ethyl acetate as eluant. The enol triflate product was obtained as a white foam and used as such in the next step. To a stirred solution of 1-((7,7-dimethyl-2-trifluoro-methanesulfonyloxy-bicyclo(2.2.1)hep-2-en-1-yl)methanesulfonyl)-4-(2-methylphenyl)-piperazine (10.5 g; 20.1 mmol) in 1:1 DMF-MeOH (150 mL) was added triethylamine (5.9 mL; 43 mmol), triphenylphosphine (317 mg; 1.21 mmol), and palladium(II)acetate (135 mg; 0.603 mmol). Carbon monoxide gas was bubbled through the solution for 15 min, and the reaction was kept under atmospheric pressure of CO for 18 h. The solvents were removed under reduced pressure and the residue was purified by pressurized silica gel column chromatography using 9:1 hexane-ethyl acetate as eluant. The title compound was obtained as a white foam from hexane.

anal: (C ₂₃ H ₃₂ N ₂ O ₄ S)			
calc.	C, 62.14;	H, 7.50;	N 6.30
found	C, 61.65;	H, 7.17;	N, 6.12

0.67 H₂O
 TLC: R_f 0.36 (1:5 EtOAc:hexanes)
 HPLC (method A): retention time 11.34 min
 FAB MS: m/z 433 (M⁺ + H)
¹H NMR (400 MHz, CDCl₃): δ 7.20 (m, 2H), 7.03 (m, 2H), 6.88 (d, J = 3 Hz, 1H), 3.72 (s, 3H), 2.33 (s, 3H), 1.09 (s, 3H), 1.01 (s, 3H)

Example 21

1-((7,7-DIMETHYL-2-CARBOXY-BICYCLO(2.2.1) HEPT-2-EN-1-YL)METHANESULFONYL)-4-(2-METHYL-PHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-methoxycarbonyl-bicyclo(2.2.1)hept-2-en-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (1.0 g; 2.3 mmol) in MeOH (10 mL) was added a solution of 4 M aqueous KOH (2.0 mL; 8.0 mmol). After 18 h, the reaction was brought to pH 1 with 5% aqueous HCl, and the solvents were removed under reduced pressure. The residue was taken up in chloroform (50 mL) and washed with water (25 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure to give the hydrochloride salt of the title compound as a white foam.

anal: (C ₂₂ H ₃₀ N ₂ O ₄ S)			
calc.	C, 57.51;	H, 6.91;	N, 6.10
found	C, 57.40;	H, 6.87;	N, 6.01

1.0 HCl, 0.25 H₂O

TLC: R_f 0.59 (92:8:0.1 CHCl₃:MeOH:HOAc)


HPLC (method A): retention time 9.77 min

FAB MS: m/z 419 (M⁺ + H)

¹H NMR (400 MHz, CD₃OD): δ 7.30 (m, 3H), 7.20 (t, 1H), 6.89 (d, J = 3 Hz, 1H), 2.43 (s, 3H), 1.11 (s, 3H), 1.00 (s, 3H)

Example 22

1-((7,7-DIMETHYL-2-(4-IMIDAZOLYL)ETHYLAMINOCARBONYL-BICYCLO(2.2.1)HEPT-2-EN-1-YL)-METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-carboxy-bicyclo(2.2.1)hept-2-en-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (100 mg; FW = 460; 0.22 mmol) in DMF (5 mL) was added histamine (30 mg; 0.27 mmol), BOP (115 mg; 0.25 mmol) and DIEA (0.12 mL; 0.69 mmol). After 18 h, the solvent was removed under

reduced pressure, the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of the title compound was obtained as a lyophilized powder.

anal: (C ₂₇ H ₃₇ N ₅ O ₃ S)			
calc.	C, 49.35;	H, 5.31;	N, 9.22
found	C, 49.25;	H, 5.39;	N, 9.20

2.1 TFA, 0.45 H₂O


HPLC (method A): retention time 8.16 min

FAB MS: m/z 512 (M⁺ + H)

¹H NMR (300 MHz, CD₃OD): δ 8.80 (s, 1H), 7.40 (s, 1H), 7.18 (m, 2H), 7.05 (d, 1H), 6.99 (t, 1H), 6.41 (d, J=3 Hz, 1H), 2.31 (s, 3H), 1.08 (s, 3H), 0.98 (s, 3H)

Example 23

1-((7,7-DIMETHYL-2-ENDO-METHOXYCARBONYL-BICYCLO (2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE



To a stirred, -78 °C solution of 1-((7,7-dimethyl-2-methoxycarbonyl-bicyclo(2.2.1)hept-2-en-1-yl)-methanesulfonyl)-4-(2-methylphenyl)piperazine (3.0 g; 6.9 mmol) in 2:1 THF-MeOH (50 mL) was added a solution of 0.1 M samarium(II) iodide in THF (250.0 mL; 25.0 mmol). After 1 h, the reaction was warmed to ambient temperature and stirred for another 1 h. The solvents were removed under reduced pressure and the residue was partitioned between ethyl acetate (100 mL) and water (50 mL). The layers were separated and the organic phase was washed with water (50 mL), dried (MgSO₄), filtered, and evaporated to dryness under reduced pressure. By ¹H NMR analysis, a 6:1 ratio of endo:exo products was obtained. The major, lower R_f isomer (endo) was obtained in pure form by pressurized silica gel column chromatography using a gradient elution of 98:2 to 95:5 hexane-ethyl acetate, followed by crystallization from ethyl acetate. The title compound was obtained as white needles.

anal: (C ₂₃ H ₃₄ N ₂ O ₄ S)			
calc.	C, 63.56;	H, 7.89;	N, 6.45
found	C, 63.31;	H, 7.83;	N, 6.43

TLC: R_f 0.44 (1:5 EtOAc:hexanes)


HPLC (method A): retention time 11.75 min

FAB MS: m/z 435 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.20 (m, 2H), 7.05 (m, 2H), 3.72 (s, 3H), 3.29 (ddd, 1H), 2.34 (s, 3H), 1.13 (s, 3H), 1.06 (s, 3H)

Example 24

1-((7,7-DIMETHYL-2-ENDO-CARBOXY-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-endo-methoxycarbonyl-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (1.0 g; 2.3 mmol) in THF (10 mL) was added a solution of 4 M aqueous NaOH (1.5 mL; 6.0 mmol). The reaction was heated to reflux for 72 h, cooled, and brought to pH 1 with 5% aqueous HCl. The solvents were removed under reduced pressure and the residue was partitioned between chloroform and water. The organic phase was separated and washed with water, dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The title compound was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The title compound was obtained as a lyophilized powder.

anal: (C ₂₂ H ₃₂ N ₂ O ₄ S)			
calc.	C, 51.92;	H, 5.99;	N, 4.94
found	C, 51.92;	H, 5.95;	N, 5.17

1.25 TFA, 0.2 H₂O

TLC: R_f 0.22 (95:5:0.5 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 10.67 min

FAB MS: m/z 421 (M⁺ + H)

¹H NMR (300 MHz, CD₃OD): δ 7.18 (m, 2H), 7.05 (d, 1H), 6.98 (t, 1H), 2.30 (s, 3H), 1.18 (s, 3H), 1.10 (s, 3H)

Example 25

1-((7,7-DIMETHYL-2-ENDO-(4-IMIDAZOLYL)ETHYLAMINOCARBONYL-BICYCLO(2.2.1)HEPTAN-1-YL)-METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-endo-carboxy-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (100 mg; 0.238 mmol) in DMF (5 mL) was histamine (35 mg; 0.32 mmol), BOP (142 mg; 0.321 mmol), and DIEA (0.13 mL; 0.75 mmol). After 18 h, the solvent was removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

anal: (C ₂₇ H ₃₉ N ₅ O ₃ S)			
calc.	C, 46.66;	H, 5.58;	N, 8.58
found	C, 46.63;	H, 5.23;	N, 8.97

2.35 TFA, 1.9 H₂O


HPLC (method A): retention time 8.99 min

FAB MS: m/z 514 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 8.40 (s, 1H), 7.1-7.3 (m, 5H), 2.39 (s, 3H), 1.05 (s, 3H), 0.98 (s, 3H)

Example 26

TWO ISOMERS OF 1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(3-METHOXYCARBONYL)-2-PYRROLIDINON-1-YL)PROPYLBICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-*exo*-hydroxy-2-*endo*-2-(1-amino)propyl-(2.2.1)bicycloheptan-1-yl)-methanesulfonyl)-4-(2-methylphenyl)piperazine (250 mg; 0.557 mmol) in methanol (3 mL) was added dimethyl itaconate (200 mg; 1.27 mmol). The reaction was heated to reflux for 18 h. The solvent was removed under reduced pressure and the residue was purified by pressurized silica gel column chromatography using 35:65 hexane-ethyl acetate as eluant. The products were obtained as white foams.

Isomer 1:

anal: (C ₃₀ H ₄₅ N ₃ O ₆ S)			
calc.	C, 62.58;	H, 7.88;	N, 7.30
found	C, 62.58;	H, 8.03;	N, 6.95

TLC: R_f 0.34 (35:65 hexane-ethyl acetate)

HPLC (method A): retention time 10.23 min

FAB MS: m/z 576 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.18 (m, 2H), 7.01 (m, 2H), 3.76 (s, 3H), 2.32 (s, 3H), 1.15 (s, 3H), 1.03 (s, 3H), 0.95 (d, J = 6 Hz, 3H)

Isomer 2:

anal: (C ₃₀ H ₄₅ N ₃ O ₆ S)			
calc.	C, 62.58;	H, 7.88;	N, 7.30
found	c, 62.43;	H, 8.07;	N, 6.95

TLC: R_f 0.23 (35:65 hexane-ethyl acetate)


HPLC (method A): retention time 10.24 min

FAB MS: m/z 576 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.20 (m, 2H), 7.03 (m, 2H), 3.74 (s, 3H), 2.32 (s, 3H), 1.15 (s, 3H), 1.03 (s, 3H), 0.95 (d, J = 6 Hz, 3H)

Example 27

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(4-PYRIDINYL)METHYLAMINO)PROPYL-BICYCLO-(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-*exo*-hydroxy-2-*endo*-2-(1-amino)propyl-(2.2.1)bicycloheptan-1-yl)-methanesulfonyl)-4-(2-methylphenyl) piperazine (50 mg; 0.11 mmol) in DMF (2 mL) was added 4-chloromethylpyridine hydrochloride (18 mg; 0.11 mmol) and potassium carbonate (50 mg; 0.36 mmol). The reaction was heated to 80°C for 18 h. The solvent was removed under reduced pressure and the residue was

purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

anal: (C ₃₀ H ₄₄ N ₄ O ₃ S)			
calc.	C, 52.07;	H, 5.89;	N, 7.06
found	C, 52.06;	H, 5.86;	N, 7.20

2.2 TFA, 0.1 H₂O

TLC: R_f 0.X


HPLC (method A): retention time 8.15 min

FAB MS: m/z 541 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 8.72 (br s, 2H), 7.85 (br s, 2H), 7.20 (m, 2H), 7.03 (m, 2H), 4.27 (AB quartet, 2H), 2.31 (s, 3H), 1.14 (s, 3H), 0.95 (overlapping s and d, 6H)

Example 28

1-((7,7-DIMETHYL-2-(3-ACETAMIDO-3,3'-DI(ETHOXYCARBONYL))PROPYLIDINE-BICYCLO(2.2.1)HEPTAN-1-YL) METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of diethyl acetamidomalonate (0.69 g; 3.2 mmol) in DMF (20 mL) was added NaH (125 mg of a 60% dispersion in mineral oil; 3.13 mmol). After 30 min, 1-((7,7-dimethyl-2-(2-chloro)ethylidene-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (0.35 g; 0.80 mmol) was added and the mixture was warmed to 50 °C for 3 h. The mixture was cooled and acetic acid (1.5 mL) was added. The solvents were removed under reduced pressure, the residue was dissolved in ethyl acetate (75 mL) and washed with water (3 x 25 mL). The organic phase was dried, filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 2:1 hexane-ethyl acetate as eluant. The title compound was obtained as a white foam.

anal: (C ₃₂ H ₄₇ N ₃ O ₇ S)			
calc.	C, 62.32;	H, 7.51;	N, 6.81
found	C, 61.96;	H, 7.71;	N, 6.55

TLC: R_f 0.36 (95:5:0.5 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 11.54 min

FAB MS: m/z 618 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.20 (m, 2H), 7.03 (m, 2H), 6.78 (s, 1H), 5.38 (br t, 1H), 4.22 (m, 4H), 2.32 (s, 3H), 2.00 (s, 3H), 1.27 (t, J = 7 Hz, 3H), 1.24 (t, J = 7 Hz, 3H), 0.97 (s, 3H), 0.78 (s, 3H)

Example 29

1-((7,7-DIMETHYL-2-(3-ACETAMIDO-3-CARBOXY) PROPYLIDINE-BICYCLO(2.2.1)HEPTAN-1-YL) METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE:



To a stirred solution of 1-((7,7-dimethyl-2-(3-acetamido-3,3'-di(ethoxycarbonyl))propylidine-(2.2.1) bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (0.10 g; 0.16 mmol) in ethanol (2 mL) was added a solution of 2 M NaOH (0.30 mL; 0.60 mmol) and the mixture was heated to reflux for 6 h. The mixture was cooled and brought to pH 2 with 5% aqueous HCl. The mixture was heated to reflux for 1 h. The solvents were removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The title compound, as a 1:1 mixture of diastereomers, was obtained as a lyophilized powder.

anal: (C ₂₇ H ₃₉ N ₃ O ₅ S)			
calc.	C, 54.37;	H, 6.53;	N, 6.56
found	C, 54.26;	H, 6.41;	N, 6.59


1.0 TFA, 0.5 H₂O

TLC: R_f 0.39 (92:8:0.1 CXCl₃:MeOH:HOAc)

HPLC (method A): retention time 9.62 min

FAB MS: m/z 518 (M⁺ + H)


¹H NMR (400 MHz, CDCl₃): δ 7.25 (m, 4H), 7.13 (m, 4H), 6.52 (d, 1H), 6.40 (d, 1H), 5.45 (m, 1H), 5.40 (m, 1H), 4.67 (m, 2H), 2.40 (s, 6H), 20.5 (s, 3H), 2.04 (s, 3H), 1.01 (s, 3H), 0.98 (s, 3H), 0.88 (s, 3H), 0.79 (s, 3H)

EXAMPLE 30**1-((7,7-DIMETHYL-2-OXO-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE**

To a stirred solution of 1-t-butyloxycarbonyl-4-(2-methylphenyl)-3-piperazinone (0.25 g; 0.86 mmol) in dichloromethane (3 mL) was added TFA (1 mL). After 1 hour the solvents were removed under reduced pressure and the residue was taken up into chloroform and evaporated several times to remove excess TFA. The residue was dissolved in chloroform (5 mL) and added to the stirred solution was 10-camphorsulfonyl chloride (376 mg; 1.50 mmol) and triethylamine (0.38 mL; 2.7 mmol). After 12 hours, the mixture was diluted with chloroform (25 mL) and extracted with 5% aqueous HCl (25 mL), water (25 mL), and aqueous NaHCO₃ (25 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 2:1 hexane-ethyl acetate as eluant. The title compound was obtained as a white foam from ether-hexane.

Anal: (C ₂₁ H ₂₈ N ₂ O ₄ S)			
Calc.	C, 62.35;	H, 6.98;	N, 6.93
Found	C, 61.78;	H, 6.98;	N, 6.82

TLC: R_f 0.30 (1:1 hexane-ethyl acetate)
 HPLC (method A): retention time 8.15 min
 FAB MS: m/z 405 (M⁺ + H)

EXAMPLE 31**1-((7,7-DIMETHYL-2-OXO-BICYCLO(2.2.1) HEPTAN-1-YL)-METHANESULFONYL)-4-(2-METHYLPHENYL)-2-METHYL-3-PIPERAZINONE**

To a stirred, 78°C solution of LDA (2.0 mmol) in THF (15 mL) was added a -78°C solution of 1-t-butyloxycarbonyl-4-(2-methylphenyl)-3-piperazinone (0.50 g; 1.7 mmol) in THF (5 mL). The resulting solution was stirred for 1 hour, when iodomethane (0.125 mL; 2.0 mmol) was added. The reaction mixture was stirred at -78°C for 30 minutes, and then the cooling bath was removed and the mixture was stirred at ambient temperature for 3 hours. Water (10 mL) and ethyl acetate (50 mL) were added. The organic layer

was separated and washed with water (25 mL) and brine (25 mL). The organic phase was dried (MgSO_4), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 85:15 hexane-ethyl acetate as eluant. The methylated product had an $R_f = 0.47$ (70:30 hexane-ethyl acetate) and an HPLC retention time of 8.32 min (Method A). The product (0.40 g; 1.3 mmol) was dissolved in chloroform (3 mL) and TFA (1 mL) was added. After 2 hours, the mixture was diluted with chloroform (50 mL) and extracted with aqueous NaHCO_3 (3 x 25 mL). The organic phase was dried (MgSO_4), filtered, and the solvent was removed under reduced pressure to give an oil (HPLC retention time 2.95 min, Method A). The residue was dissolved in chloroform (20 mL) and to the stirred solution was added 10-camphorsulfonyl chloride (0.41 g; 1.6 mmol) and triethylamine (0.28 mL; 2.0 mmol). After 12 hours, the mixture was diluted with chloroform (25 mL) and extracted with 5% aqueous HCl (25 mL), water (25 mL), and aqueous NaHCO_3 (2 x 25 mL). The organic phase was dried (MgSO_4), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 2:1 hexane-ethyl acetate as eluant. The title compound, as a 1:1 mixture of diastereomers, was obtained as a white solid from hexane-ether.

Anal: ($\text{C}_{22}\text{H}_{30}\text{N}_2\text{O}_4\text{S}$)			
Calc.	C, 63.13;	H, 7.23;	N, 6.69
Found	C, 63.46;	H, 7.09;	N, 6.74

TLC: R_f 0.27 (60:40 hexane-ethyl acetate)


HPLC (method A): retention time 8.52 min

FAB MS: m/z 419 ($\text{M}^+ + \text{H}$)

^1H NMR (300 MHz, CDCl_3): δ 7.1-7.3 (m, 8H), 4.62 (overlapping quartets, 2H), 2.21 (s, 3H), 2.20 (s, 3H), 1.68 (overlapping doublets, 6H), 1.13 (s, 3H), 1.11 (s, 3H), 0.91 (s, 3H), 0.89 (s, 3H)

EXAMPLE 32

1-((7,7-DIMETHYL-2-EXO-HYDROXY-BICYCLO (2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYL-PHENYL)-2-METHYL-PIPERAZINE



To a stirred, 0°C solution of 1-((7,7-dimethyl-2-oxo-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)-2-methyl-3-piperazinone (0.15 g; 0.36 mmol) in THF (5 mL) was added a 1.0 M solution of LAH in THF (1.1 mL; 1.1 mmol). The resulting solution was warmed to ambient temperature and stirred for 3 hours. The reaction was quenched by adding aqueous NaOH to give a white precipitate. The mixture was diluted with ethyl acetate and the solids were removed by filtration through Celite. The filtrate solvents were removed under reduced pressure and the residue was purified by pressurized silica gel column chromatography using 9:1 hexaneethyl acetate as eluant to give 1-((7,7-dimethyl-2-exo-hydroxy-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)-2-methyl-2,3-dehydro-piperazine (FAB MS: m/z 405 ($\text{M}^+ + \text{H}$); olefinic proton at 5.8 ppm in the ^1H NMR spectrum). This product (75 mg; 0.19 mmol) was dissolved in triethylsilane (2 mL) and to the stirred solution was added TFA (0.030 mL; 0.38 mmol). After 18 hours, the solvents were removed under reduced pressure and the residue was dissolved in ethyl acetate (20 mL) and washed with aqueous NaHCO_3 (2 x 10 mL). The organic phase was dried (MgSO_4), filtered, and the solvent was removed under reduced pressure. The residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The title compound, as a 1:1 mixture of diastereomers,

was obtained as a lyophilized powder.

Anal: (C ₂₂ H ₃₄ N ₂ O ₃ S)			
Calc.	C, 63.13;	H, 7.23;	N, 6.69
Found	C, 63.46;	H, 7.09;	N, 6.74

TLC: R_f 0.27 (60:40 hexane-ethyl acetate)


HPLC (method A): retention time 14.33 min

FAB MS: m/z 407 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.20 (m, 4H), 7.06 (m, 4H), 4.20 (m, 2H), 2.36 (s, 6H), 1.55 (overlapping doublets, 6H), 1.09 (s, 6H), 0.86 (s, 6H)

EXAMPLE 33

1-((7,7-DIMETHYL-2-OXIMINO-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE



To a stirred solution of 1-((7,7-dimethyl-2-oxo-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (65.0 g; 166 mmol) in pyridine (250 mL) was added hydroxylamine hydrochloride (35.0 g; 0.504 mol). The solution was heated to 70 °C for 18 h. The solvent was removed under reduced pressure, the residue was taken up in chloroform (500 mL) and washed with aqueous NaHCO₃ (2 x 200 mL), water (100 mL), and 5% aqueous HCl (2 x 200 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The title compound crystallized from ethyl acetate, giving off-white needles (57 g; 84%), mp 174-175°C.


Anal: (C ₂₁ H ₃₁ N ₃ O ₃ S)			
Calc.	C, 62.19;	H, 7.71;	N, 10.36
Found	C, 62.29;	H, 7.63;	N, 10.15

TLC: R_f 0.40 (75:25 hexane-ethyl acetate)

HPLC (method A): retention time 9.98 min


FAB MS: m/z 406 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.90 (br s, 1H), 7.18 (m, 2H), 7.02 (m, 2H), 3.47 (m, 4H), 4.43 (d, J = 14.4 Hz, 1H), 3.00 (m, 4H), 2.92 (d, J = 14.4 Hz, 1H), 2.4-2.6 (m, 2H), 2.31 (s, 3H), 2.09 (d, J = 16.9 Hz, 1H), 1.95 (m, 2H), 1.80 (m, 1H), 1.32 (m, 1H), 1.08 (s, 3H), 0.87 (s, 3H)

EXAMPLE 34**1-((7,7-DIMETHYL-2-ENDO-AMINO-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE**

To a stirred solution of 1-((7,7-dimethyl-2-oximino-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (35.0 g; 86 mmol) in 2-methoxyethanol (500 mL) containing Raney Nickel alloy (105.0 g) was added sodium hydroxide solution (17.2 g; 430 mmol dissolved in 75 mL) dropwise over 30 min. During the addition heat and gas was evolved. The mixture was stirred at ambient temperature for 16 h, at which time TLC indicated complete consumption of starting oxime and a ca. 4:1 mixture of endo (lower R_f and exo (higher R_f) amine products. The mixture was filtered through Celite and the filter-cake was washed with methanol and ethyl acetate. The solvents were removed under reduced pressure and the resulting solid was dispersed in water and filtered. The dried solid was purified by pressurized silica gel column chromatography, using a 93:3 to 94:6 A:B gradient elution (A=chloroform, B=5% NH₄OH/MeOH). The title compound was obtained as a white foam (24 g; 70%).

FAB MS: m/z 392 (M⁺ + H)

EXAMPLE 35**1-((7,7-DIMETHYL-2-ENDO-(2S-(TERT-BUTYLOXYCARBONYLAMINO)-4-(METHYLSULFONYL)-BUTYRAMIDO)-BICYCLO(2.2.1)-HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE**

To a stirred solution of 1-((7,7-dimethyl-2-endo-amino-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (2.0 g; 5.1 mmol) in DMF (20 mL) was added Na-Boc-L-methionine sulfonate (1.5 g;


5.3 mmol), BOP reagent (2.5 g; 5.6 mmol), followed by DIEA (1.85 mL; 10.6 mmol). After being stirred at ambient temperature for 1 h, more DIEA (ca. 0.1 mL) was added to obtain a pH 8 solution. The solution was stirred for another 1 h, when the solvent was removed under reduced pressure. The residue was dissolved in EtOAc (150 mL) and washed with 5% aqueous HCl (2 x 50 mL), water (2 x 50 mL), and aqueous NaHCO₃ (2 x 75 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography, using 4:1 EtOAc-hexanes as eluant. The title compound was obtained as a solid from methanol (2.8 g; 85%).

Anal: (C ₃₁ H ₅₀ N ₄ O ₇ S ₂)				
Calc.	C, 55.78;	H, 7.76;	N, 8.39	0.7 H ₂ O
Found	C, 55.57;	H, 7.70;	N, 8.36	

TLC: R_f 0.73 (95:5 CHCl₃:MeOH)
 HPLC (method A): retention time 11.02 min
 FAB MS: m/z 655 (M⁺ + H)
¹H NMR (300 MHz, CDCl₃): δ 7.19 (m, 2H), 7.04 (m, 2H), 5.38 (br d, 1H), 4.32 (q, J = 7.4 Hz, 1H), 4.22 (m, 1H), 2.94 (s, 3H), 2.32 (s, 3H), 1.45 (s, 9H), 1.00 (s, 3H), 0.98 (s, 3H)

EXAMPLE 36

1-((7,7-DIMETHYL-2-ENDO-(2S-AMINO-4-(METHYLSULFONYL)-BUTYRAMIDO)-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE



To a stirred solution of 1-((7,7-dimethyl-2-endo-(2S-tert-butyloxycarbonylamino-4-(methylsulfonyl)butyramido)-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (2.5 g; 3.8 mmol) in dichloromethane (15 mL) was added TFA (5 mL). After 1 h, the solvents were removed under reduced pressure. The residue was dissolved in chloroform (100 mL) and washed with aqueous NaHCO₃ (2 x 75 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 95:5:0.5 CHCl₃:MeOH:NH₄OH as eluant. The title compound was obtained as a white foam from EtOAc (1.9 g; 90%).

Anal: (C ₂₆ H ₄₂ N ₄ O ₅ S ₂)				
Calc.	C, 56.14;	H, 7.75;	N, 9.29	0.55 EtOAc
Found	C, 55.94;	H, 7.74;	N, 9.31	

TLC: R_f 0.17 (95:5:0.5 CHCl_3 :MeOH:NH₄OH)


HPLC (method A): retention time 8.50 min

FAB MS: m/z 455 ($M^+ + H$)

¹H NMR (300 MHz, CDCl_3): δ 7.67 (d, J = 8.4 Hz, 1H), 7.20 (m, 2H), 7.02 (m, 2H), 4.43 (m, 1H), 2.94 (s, 3H), 2.31 (s, 3H), 1.03 (s, 3H), 0.97 (s, 3H)

EXAMPLE 37

1-((7,7-DIMETHYL-2-ENDO-(2S-(IMIDAZOL-4-YLACETYLAMINO)-4-(METHYLSULFONYL)BUTYRAMIDO)-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE



To a stirred solution of 1-((7,7-dimethyl-2-endo-(2S-amino-4-(methylsulfonyl)butyramido)-bicyclo(2.2.1)-heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl) piperazine (250 mg; 0.45 mmol) in DMF (5 mL) was added 4-imidazole acetic acid hydrochloride (110 mg; 0.68 mmol), BOP (265 mg; 0.60 mmol), and DIEA (0.355 mL; 2.0 mmol). The solution was stirred at ambient temperature for 18 h. The solvent was removed under reduced pressure, and the residue was suspended in EtOAc (100 mL) and filtered through Celite to remove red polymer. The filtrate was washed with 5% aqueous HCl (50 mL), water (50 mL), and aqueous NaHCO₃ (2 x 50 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 92:8:0.8 CHCl_3 :MeOH:NH₄OH as eluant. The title compound was obtained as a solid from EtOAc (230 mg; 78%).

Anal: ($\text{C}_{31}\text{H}_{46}\text{N}_6\text{O}_6\text{S}_2$)					
Calc.	C, 53.74;	H, 7.32;	N, 11.26	0.6 EtOAc,	1.7H ₂ O
Found	C, 53.74;	H, 7.00;	N, 11.25		

TLC: R_f 0.22 (90:10:0.5 CHCl_3 :MeOH:NH₄OH)


HPLC (method A): retention time 8.49 min

FAB MS: m/z 663 ($M^+ + H$)

¹H NMR (300 MHz, CDCl_3): δ 7.73 (overlapping singlet and broad singlet, 2H), 7.38 (br d, 1H), 7.18 (m, 2H), 7.02 (m, 2H), 6.96 (s, 1H), 4.68 (br q, J = ca. 5 Hz, 1H), 4.27 (m, 1H), 3.62 (br s, 2H), 2.92 (s, 3H), 2.30 (s, 3H), 1.00 (s, 3H), 0.98 (s, 3H)

EXAMPLE 38

1-((7,7-DIMETHYL-2-ENDO-(2S-(DIMETHYLAMINO)-4-(METHYLSULFONYL)BUTYRAMIDO)-BICYCLO-(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE



To a stirred solution of 1-((7,7-dimethyl-2-endo-(2S-amino-4-(methylsulfonyl)butyramido)-bicyclo(2.2.1)-heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl) piperazine (250 mg; 0.45 mmol) in 1:1 HOAc:MeOH (10 mL) was added 37% aqueous formaldehyde (2 mL) and NaBH₃CN (60 mg; 0.95 mmol). The solution was stirred at ambient temperature for 4 h. Aqueous NaHCO₃ (2 mL) was added and the solvents were removed under reduced pressure. The residue was suspended in EtOAc (75 mL) and washed with water (2 x 50 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The title compound was obtained as a white foam from EtOAc (190 mg; 72%).

Anal: (C₂₈H₄₆N₄O₅S₂)

Calc.	C, 57.56;	H, 8.01;	N, 9.20	0.3 EtOAc,
Found	C, 57.41;	H, 7.98;	N, 9.20	

TLC: R_f 0.26 (95:5:0.5 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 9.10 min

FAB MS: m/z 583 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.62 (br s, 1H), 7.18 (m, 2H), 7.02 (M, 2H), 4.37 (m, 1H), 2.92 (s, 3H), 2.36 (s, 6H), 2.30 (s, 3H), 1.02 (s, 3H), 0.98 (s, 3H)

EXAMPLE 39

1-((7,7-DIMETHYL-2-ENDO-BENZYLOXYCARBONYLAMINO-BICYCLO(2.2.1)HEPTAN-1-YL)-METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE



To a 0°C stirred solution of 1-((7,7-dimethyl-2-endo-amino-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (1.20 g; 3.07 mmol) in CHCl₃ (100 mL) was added DIEA (0.80 mL; 4.6 mmol) and benzyl chloroformate (0.58 g; 3.4 mmol). The solution was stirred at 0°C for 1 h and then at ambient temperature for 4 h. The reaction mixture was washed with 5% aqueous HCl (2 x 50 mL) and aqueous NaHCO₃ (100 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 1:4 EtOAc-hexanes as eluant. The title compound was obtained as a white foam. (1.45 g; 90%).

Anal: (C ₂₉ H ₃₉ N ₃ O ₄ S)					
Calc.	C, 65.75;	H, 7.53;	N, 7.77	0.15 EtOAc,	0.1 H ₂ O
Found	C, 65.90;	H, 7.49;	N, 7.80		


TLC: R_f 0.38 (1:3 EtOAc:hexanes)

HPLC (method A): retention time 12.18 min

FAB MS: m/z 526 (M⁺ + H)

EXAMPLE 40

1-((7,7-DIMETHYL-2-ENDO-METHYL(BENZYLOXYCARBONYLAMINO-BICYCLO(2.2.1)HEPTAN-1-YL)-METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE



To a 0°C stirred solution of 1-((7,7-dimethyl-2-endo-benzyloxycarbonylamino-bicyclo(2.2.1)heptan-1-yl)-methanesulfonyl)-4-(2-methylphenyl)piperazine (1.46 g; 2.78 mmol) in DMF (20 mL) was added iodomethane (0.435 mL; 7.00 mmol) and sodium hydride (0.139 mg of a 60% dispersion in mineral oil; 3.48 mmol). The solution was stirred at 0°C for 1 h and then at ambient temperature for 18 h. The reaction mixture was treated with HOAc (1 mL) and the solvents were removed under reduced pressure. The residue was dissolved in EtOAc (100 mL) and washed with aqueous NaHCO₃ (2 x 50 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 1:5 EtOAc-hexanes as eluant. The title compound was obtained as a white foam. (1.40 g; 93%).

Anal: (C ₃₀ H ₄₁ N ₃ O ₄ S)				
Calc.	C, 66.03;	H, 7.70;	N, 7.70	0.33 H ₂ O
Found	C, 66.03;	H, 7.63;	N, 7.68	

TLC: R_f 0.44 (1:4 EtOAc:hexanes)

HPLC (method A): retention time 12.86 min

FAB MS: m/z 540 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.25-7.45 (m, 5H), 7.20 (m, 2H), 7.02 (m, 2H), 5.11 (AB quartet, 2H), 4.83 (m, 1H), 3.03 (s, 3H), 2.32 (s, 3H), 1.04 (s, 3H), 0.96 (s, 3H)

EXAMPLE 41

1-((7,7-DIMETHYL-2-ENDO-METHYL(2S-AMINO-4-(METHYLSULFONYL)BUTANOYL)AMINO-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE

To a stirred, argon purged solution of 1-((7,7-dimethyl-2-endo-methyl(benzyloxycarbonylamino-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (1.1 g; 2.0 mmol) in 96:4 MeOH-HCO₂H (25 mL) was added palladium black (0.4 g). The reaction mixture was stirred for 16 h at ambient temperature. The catalyst was removed by filtration through Celite, and the filtrate solvents were removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 95:5:0.5 CHCl₃:MeOH:NH₄OH as eluant. The product, 1-((7,7-dimethyl-2-endo-methylamino-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine, was obtained as a white foam. (0.79 g; 95%). To a stirred solution of 1-((7,7-dimethyl-2-endo-methylamino-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (0.700 g; 1.73 mmol) in CHCl₃ (60 mL) was added the acid fluoride of Na-Fmoc-L-methionine sulfone (1.23 g; 3.03 mmol) and DIEA (0.52 mL; 3.0 mmol). The mixture was stirred at ambient temperature for 24 h, and then extracted with 5% aqueous HCl (30 mL), water (30 mL), and aqueous

NaHCO₃ (2 x 30 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was dissolved in DMF (10 mL), and to this solution was added diethylamine (2 mL). The mixture was stirred at ambient temperature for 6 h. The solvents were removed under reduced pressure and the residue was purified by pressurized silica gel column chromatography using 95:5:0.5 CHCl₃:MeOH:NH₄OH as eluant. The title compound was obtained as a foam from CHCl₃-ether (0.71 g; 61%).

Anal: (C ₂₇ H ₄₄ N ₄ O ₅ S ₂)					
Calc.	C, 56.26;	H, 7.80;	N, 9.40	0.1 CHCl ₃ ,	0.2 ether
Found	C, 56.21;	H, 7.79;	N, 9.22		

TLC: R_f 0.10 (95:5:0.5 CHCl₃:MeOH:NH₄OH)

HPLC (method A): retention time 9.01 min

FAB MS: m/z 569 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.18 (m, 2H), 7.03 (m, 2H), 5.20 (ddd, 1H), 3.95 (dd, J = 9.3, 4.1 Hz, 1H), 3.18 (s, 3H), 2.91 (s, 3H), 2.30 (s, 3H), 1.06 (s, 3H), 0.96 (s, 3H)

EXAMPLE 42

1-((7,7-DIMETHYL-2-ENDO-METHYL(2S-DIMETHYLAMINO-4-(METHYLSULFONYL)BUTANOYL)AMINO-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE

To a stirred solution of 1-((7,7-dimethyl-2-endo-methyl(2S-amino-4-(methylsulfonyl)butanoyl)amino-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (150 mg; 0.264 mmol) in 1:1 HOAc:MeOH (6 mL) was added 37% aqueous formaldehyde (1 mL) and NaBH₃CN (30 mg; 0.47 mmol). The solution was stirred at ambient temperature for 4 h. Aqueous NaHCO₃ (1 mL) was added and the solvents were removed under reduced pressure. The residue was suspended in EtOAc (50 mL) and washed with water (2 x 25 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by preparative reverse phase HPLC using a water-acetonitrile gradient containing 0.1% TFA. The TFA salt of the title compound was obtained as a lyophilized powder.

Anal: (C ₂₉ H ₄₈ N ₄ O ₅ S ₂)					
Calc.	C, 44.88;	H, 5.94;	N, 6.16	2.5 TFA,	1.5 H ₂ O
Found	C, 44.80;	H, 5.94;	N, 6.18		

TLC: R_f 0.45 (95:5:0.5 CHCl₃:MeOH:NH₄OH)

HPLC (method A): retention time 9.04 min

FAB MS: m/z 597 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.2-7.3 (m, 4H), 5.15 (m, 1H), 4.79 (br t, 1H), 3.21 (s, 3H), 2.98 (s, 3H), 2.95 (s, 6H), 2.43 (s, 3H), 1.07 (s, 3H), 0.97 (s, 3H)

EXAMPLE 43

1-((7,7-DIMETHYL-2-ENDO-(4-IMIDAZOLYL)ACETYL)AMINO-BICYCLO(2.2.1)HEPTAN-1-YL)-METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINONE

To a stirred solution of 1-((7,7-dimethyl-2-endo-amino-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (1.50 g; 3.84 mmol) in DMF (30 mL) was added 4-imidazole acetic acid hydrochloride (0.938 g; 5.76 mmol), BOP (2.13 g; 4.80 mmol), and DIEA (2.61 mL; 15.0 mmol). The reaction mixture was stirred for 24 h at ambient temperature, and the solvent was removed under reduced pressure. The residue was suspended in EtOAc (100 mL) and filtered through Celite to remove dried polymer. The filtrate was washed with aqueous NaHCO₃ (2 x 50 mL) and water (2 x 50 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography using 92:8:0.8 CHCl₃:MeOH:NH₄OH as eluant. The title compound was obtained as white foam.

FAB MS: m/z 500 ($M^+ + H$)EXAMPLE 445 1-((7,7-DIMETHYL-2-ENDO-(2-(4-IMIDAZOLYL)PROPANOYL)AMINO-BICYCLO(2.2.1)HEPTAN-1-YL)-METHANESULFONYL)-4-(2-METHYLPHENYL)-3-PIPERAZINE

To a stirred solution of 1-((7,7-dimethyl-2-endo-amino-bicyclo(2.2.1)heptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (1.1 g; 2.8 mmol) in DMF (25 mL) was added 2-(1-benzyloxymethyl-5-imidazolyl)-
 10 propionic acid hydrochloride (0.920 g; 3.10 mmol), BOP (1.35 g; 3.05 mmol), and DIEA (1.50 mL; 8.61 mmol). The reaction mixture was stirred for 1 h at ambient temperature, and more DIEA (ca. 0.2 mL) was added to bring the mixture to pH 8. After another 1 h, the solvent was removed under reduced pressure. The residue was dissolved in $CHCl_3$ (150 mL) and washed with aqueous $NaHCO_3$ (2 x 50 mL) and water (2 x 50 mL). The organic phase was dried ($MgSO_4$), filtered, and the solvent was removed under reduced
 15 pressure to give a solid. Recrystallization from EtOAc gave crystals (0.51 g) which, by 1H NMR analysis, proved to be a 90:10 mixture of isomers (product A). The filtrate was purified by pressurized silica gel column chromatography using 95:5 $CHCl_3$:MeOH as eluant, giving a white foam (1.0 g). 1H NMR indicated this material to be a 1:2 mixture of isomers (product B). Products A and B were individually deblocked by hydrogenation for 24 h at ambient temperature in 3:1 MeOH:HOAc using 25 weight% palladium black under
 20 1 atmosphere of hydrogen. The catalyst was removed by filtration through Celite and the solvents were removed under reduced pressure. The residue derived from product A was purified by preparative reverse phase HPLC using a water-acetonitrile gradient containing 0.1% TFA. The TFA salt of the title compound (90:10 mixture by 1H NMR) was obtained as a lyophilized powder. Product B was purified by
 25 pressurized silica gel column chromatography using 95:5:0.5 $CHCl_3$:MeOH: NH_4OH as eluant. The title compound was obtained as white foam from $CHCl_3$ -ether (1:2 mixture by 1H NMR). The two isomers had identical chromatographic behavior.

30

Anal: ($C_{27}H_{37}N_5O_3S$)					
Calc.	C, 60.36;	H, 7.49;	N, 12.46	0.25 $CHCl_3$,	0.25 ether
Found	C, 60.49;	H, 7.26;	N, 12.48		

35 TLC: R_f 0.30 (93:7:0.7 $CHCl_3$:MeOH: NH_4OH)

HPLC (method A): retention time 8.79 min

FAB MS: m/z 514 ($M^+ + H$) 1H NMR (400 MHz, $CDCl_3$): δ 7.75 (br s, 1H), 7.20 (m, 2H), 7.0 (m, 3H), 4.40 (m, 1H), 2.30, 2.29 (two singlets, ca. 2:1 ratio, 3H), 1.57, 1.53 (two doublets, $J=7$ Hz, ca. 2:1 ratio, 3H), 1.00 (s, 3H), 0.96 (s, 3H)

40 L-369,076

45

Anal: ($C_{27}H_{37}N_5O_3S$)				
Calc.	C, 48.91;	H, 5.36;	N, 9.03	2.3 TFA
Found	C, 48.99;	H, 5.21;	N, 9.03	

TLC: R_f 0.30 (93:7:0.7 $CHCl_3$:MeOH: NH_4OH)

HPLC (method A): retention time 8.79 min

FAB MS: m/z 514 ($M^+ + H$)50 1H NMR (400 MHz, $CDCl_3$): δ 8.43 (s, 1H), 7.70 (d, 1H), 7.25 (m, 2H), 7.20 (s, 1H), 7.15 (m, 2H), 4.40 (m, 1H), 4.03 (q, $J=7$ Hz, 1H), 2.38 (s, 3H), 1.57 (d, $J=7$ Hz, 3H), 1.00 (s, 3H), 0.95 (s, 3H)

55

EXAMPLE 45


1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(L-N-(METHOXYCARBONYLETHYL)PROLYL)AMINO)-
PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE

5

10

15

20



25 To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(L-prolyl)amino)propyl(2.2.1)-
 bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (1.50 g; 2.74 mmol) in methanol (15 mL)
 was added methyl acrylate (0.310 mL; 3.43 mmol). After 72 h at ambient temperature, the solvent was
 removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an
 acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a
 30 lyophilized powder.

35

Anal: (C ₃₃ H ₅₂ N ₄ O ₆ S)				
Calc.	C, 53.10;	H, 6.59;	N, 6.82	1.65 TFA
Found	C, 53.09;	H, 6.58;	N, 6.88	

TLC: R_f 0.55 (95:5 CHCl₃:MeOH)

HPLC (method A): retention time 9.45 min

FAB MS: m/z 633 (M⁺ + H)

40

¹H NMR (400 MHz, CDCl₃): δ 7.18 (m, 2H), 7.03 (m, 2H), 4.55 (m, 1H), 3.72 (s, 3H), 2.32 (s, 3H), 1.15 (s,
 3H), 1.04 (s, 3H), 1.01 (d, J = 6 Hz, 3H)


45

50

55

EXAMPLE 46

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(L-N-(CARBOXYETHYL)PROLYL)AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(L-N-(methoxycarbonyl)ethyl)prolyl)-amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (1.00 g; FW = 821; 1.22 mmol) in THF (15 mL) was added 1 M NaOH until a pH 10 solution persisted for 1 h. The solution was evaporated under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

Anal: (C ₃₂ H ₅₀ N ₄ O ₆ S)				
Calc.	C, 51.88;	H, 6.34;	N, 6.80	1.8 TFA
Found	C, 51.87;	H, 6.28;	N, 6.82	

TLC: R_f 0.40 (80:20:2 CHCl₃:MeOH:NH₄OH)

HPLC (method A): retention time 8.88 min

FAB MS: m/z 619 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 8.50 (br s 1H), 7.20 (m, 2H), 7.05 (m, 2H), 2.33 (s, 3H), 1.12 (s, 3H), 1.03 (s, 3H), 0.99 (d, J = 6 Hz, 3H)

EXAMPLE 47


1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(3-PIPERIDINYLCARBONYL)AMINO)PROPYL-BICYCLO-(2.2.1)HEPTAN-1-YL) METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE

5

10

15

20



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (2.50 g; 5.57 mmol) in DMF (35 mL) was added N-Fmoc-piperidine-3-carboxylic acid (2.15 g; 6.13 mmol), BOP (2.75 g; 6.20 mmol), and DIEA (2.16 mL; 12.4 mmol). After 16 h, diethylamine (6 mL) was added and the solution was stirred at ambient temperature for 4 h. The solvents were removed under reduced pressure and the residue was dissolved in EtOAc (150 mL) and washed with aqueous NaHCO₃ (2 x 75 mL) and water (2 x 75 mL). The organic phase was dried (MgSO₄), filtered, and the solvent was removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography, using 93:7:0.7 CHCl₃:MeOH:NH₄OH as eluant. The title compound (1:1 mixture of diastereomers) was obtained as a white foam.

35

Anal: (C ₃₀ H ₄₈ N ₄ O ₄ S)				
Calc.	C, 56.37;	H, 7.49;	N, 8.54	0.8 CHCl ₃
Found	C, 56.49;	H, 7.44;	N, 8.50	

40

TLC: R_f 0.40 (90:10:1 CHCl₃:MeOH:NH₄OH)

HPLC (method A): retention time 8.67 min

FAB MS: m/z 561 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 7.50 (br s, 1H), 7.20 (m, 2H), 7.02 (m, 2H), 2.30 (s, 3H), 1.17 (s, 3H), 1.00-1.04 (overlapping singlet and doublet, 6H)

45

EXAMPLE 48

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(3-(1-METHOXYCARBONYLETHYL)-PIPERIDINYLCARBONYL)AMINO)PROPYL-BICYCLO-(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE

50

55



25

30

TLC: R_f 0.35 (95:5 CHCl₃:MeOH)

HPLC (method A): retention time 10.71 min

FAB MS: m/z 647 ($M^+ + H$)

35


EXAMPLE 49

40

45

50

55



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(3-(1-methoxycarbonyl)-
 20 piperidinylcarbonyl)amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine
 (0.30 g; 0.46 mmol) in THF (10 mL) was added 1 M NaOH until a pH 10 solution persisted for 1 h. The
 solution was evaporated under reduced pressure and the residue was purified by preparative reverse phase
 HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound (1:1 mixture
 25 of diastereomers) was obtained as a lyophilized powder.

30

Anal: (C ₃₃ H ₅₂ N ₄ O ₆ S)					
Calc.	C, 51.59;	H, 6.44;	N, 6.54	1.9 TFA,	0.4 H ₂ O
Found	C, 51.60;	H, 6.44;	N, 6.83		

TLC: R_f 0.15 (80:20:2 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 10.27 min

FAB MS: m/z 633 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.20 (m, 2H), 7.05 (m, 2H), 2.39, 2.32 (two singlets, 3H), 1.12, 1.11 (two
 35 singlets, 3H), 0.95-1.03 (two coincident singlets and two overlapping doublets, 6H)

EXAMPLE 50

40 1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(3-(1-ETHOXYCARBONYLMETHYL)-
PIPERIDINYLCARBONYL)AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-
METHYLPHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(3-piperidinylcarbonyl)amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (0.50 g; 0.89 mmol) in DMF (5 mL) was added ethyl bromoacetate (0.110 mL; 0.99 mmol) and DIEA (0.172 mL; 0.99 mmol). After 24 h at ambient temperature, the solvent was removed under reduced pressure and the residue was dissolved in EtOAc (50 mL) and washed with 5% aqueous citric acid (25 mL), water (25 mL), and aqueous NaHCO₃ (25 mL). The organic phase was dried (MgSO₄), filtered, and the solvents were removed under reduced pressure. The residue was purified by pressurized silica gel column chromatography, using 1:1 EtOAc:CHCl₃ as eluant. The title compound (1:1 mixture of diastereomers) was obtained as a white foam.

Anal: (C₃₄H₅₄N₄O₆S)

Calc.	C, 58.66;	H, 7.77;	N, 7.93	0.5 CHCl ₃
Found	C, 58.87;	H, 7.83;	N, 7.88	

TLC: R_f 0.28 (1:1 CHCl₃:EtOAc)


HPLC (method A): retention time 9.76 min

FAB MS: m/z 647 (M⁺ + H)

¹H NMR (300 MHz, CDCl₃): δ 8.2 (very br s, 1H), 7.18 (m, 2H), 7.03 (m, 2H), 4.20 (two very closely spaced quartets, 2H), 2.30, 2.31 (two singlets, 3H), 1.28 (t, J = 7 Hz, 3H), 1.07, 1.08 (two singlets, 3H), 1.03-1.08 (two coincident singlets and two overlapping doublets, 6H)

EXAMPLE 51

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(3-(1-CARBOXYMETHYL)PIPERIDINYL-CARBONYL)-AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(3-(1-methoxycarbonyl)piperidinylcarbonyl)amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (0.360 g; 0.555 mmol) in THF (5 mL) was added 1 M NaOH until a pH 10 solution persisted for 1 h. The solution was made acidic by the addition of HOAc (1 mL) and evaporated under reduced pressure. The residue was suspended in CH₂Cl₂ and filtered. The filtrate was evaporated under reduced pressure several times from CH₂Cl₂ to give the title compound (1:1 mixture of diastereomers) as a white foam.

Anal: (C₃₂H₅₀N₄O₆S)

Calc.	C, 58.27;	H, 7.62;	N, 7.99	1.0 NaOAc
Found	C, 58.47;	H, 7.71;	N, 7.90	

TLC: R_f 0.55 (85:15 CHCl₃:MeOH)

HPLC (method A): retention time 8.77 min

FAB MS: m/z 619 ($M^+ + H$)

1H NMR (300 MHz, CD_3OD): δ 7.15 (m, 2H), 7.05 (d, $J=7.3$ Hz, 1H), 6.96 (t, $J=7.3$ Hz, 1H), 2.31 (s, 3H), 1.17 (s, 3H), 1.03 (s, 3H), 0.98 (d, $J=6$ Hz, 3H)


5 EXAMPLE 52

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(L-N-(ETHOXYCARBOXYMETHYL)PROLYL)AMINO)-
PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL) PIPERAZINE

10

15

20



25

30

To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(L-prolyl)amino)propyl(2.2.1)-bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (0.20 g; 0.37 mmol) in DMF (5 mL) was added ethyl bromoacetate (0.045 mL; 0.40 mmol) and DIEA (0.071 mL; 0.41 mmol). After 24 h at ambient temperature, the solvent was removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

35

Anal: ($C_{33}H_{52}N_4O_5S$)

Calc.	C, 54.25;	H, 6.79;	N, 7.07	1.4 TFA
Found	C, 54.25;	H, 6.78;	N, 7.02	

40

TLC: R_f 0.50 (1:1 EtOAc:CHCl₃)

HPLC (method A): retention time 9.68 min

FAB MS: m/z 633 ($M^+ + H$)

1H NMR (400 MHz, CD_3OD): δ 7.17 (m, 2H), 7.06 (d, $J=6$ Hz, 1H), 6.98 (t, $J=6$ Hz, 1H), 4.25 (m, 3H), 4.08 (d, $J=15$ Hz, 1H), 2.32 (s, 3H), 1.27 (t, $J=7$ Hz, 3H), 1.18 (s, 3H), 1.03 (s, 3H), 1.01 (d, $J=6$ Hz, 3H)


45

50

55

EXAMPLE 53

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(L-N-(CARBOXYMETHYL)PROYL)AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(L-(N-ethoxycarbonylmethyl)proyl)-amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (0.20 g; 0.32 mmol) in THF (5 mL) was added 1 M NaOH until a pH 10 solution persisted for 1 h. The solvent was removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

Anal: (C₃₁H₄₈N₄O₆S)

Calc.	C, 52.64;	H, 6.43;	N, 7.22	1.5 TFA
Found	C, 52.49;	H, 6.51;	N, 7.22	

TLC: R_f 0.40 (80:20:2 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 8.79 min

FAB MS: m/z 605 (M⁺ + H)

¹H NMR (400 MHz, CD₃OD): δ 7.17 (m, 2H), 7.07 (d J=5 Hz, 1H), 6.99 (t, J=5 Hz, 1H), 4.30 (dd, J=4, 5 Hz, 1H), 4.21 (d, J=14 Hz, 1H), 4.04 (d, J=14 Hz, 1H), 2.32 (s, 3H), 1.18 (s, 3H), 1.03 (s, 3H), 1.01 (d, J=7 Hz, 3H)

EXAMPLE 54

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(4-PIPERIDINYLCARBONYL)AMINO)PROPYL-BICYCLO-(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (1.50 g; 3.34 mmol) in DMF (20 mL) was added N-Fmoc-piperidine-4-carboxylic acid (1.29 g; 3.67 mmol), BOP (1.64 g; 3.70 mmol), and DIEA (1.28 mL; 7.34 mmol). After 16 h, diethylamine (5 mL) was added and the solution was stirred at ambient temperature for 4 h. The solvents were removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

Anal: (C ₃₀ H ₄₈ N ₄ O ₄ S)					
Calc.	C, 51.93;	H, 6.43;	N, 7.15	1.95 TFA,	0.05 H ₂ O
Found	C, 51.93;	H, 6.36;	N, 7.28		

TLC: R_f 0.15 (90:10:1 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 8.33 min

FAB MS: m/z 561 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.20 (m, 3H), 7.08 (m, 2H), 2.33 (s, 3H), 1.14 (s, 3H), 1.02 (s, 3H), 1.00 (d, J = 6 Hz, 3H)

EXAMPLE 55

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(4-(1-METHOXYCARBONYLETHYL)-PIPERIDINYLCARBONYL)AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(4-(2-methylphenyl)piperidinyldicarbonyl)amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (0.30 g; 0.53 mmol) in methanol (5 mL) was added methyl acrylate (0.072 mL; 0.80 mmol). After 48 h at ambient temperature, the solvent was removed under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

Anal: (C ₃₄ H ₅₄ N ₄ O ₆ S)					
Calc.	C, 53.04;	H, 6.65;	N, 6.60	1.75 TFA,	0.15 H ₂ O
Found	C, 53.05;	H, 6.62;	N, 6.69		

TLC: R_f 0.25 (95:5 CHCl₃:MeOH)


HPLC (method A): retention time 9.02 min

FAB MS: m/z 647 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.45 (br t, 1H), 7.21 (m, 2H), 7.09 (m, 2H), 3.72 (s, 3H), 2.33 (s, 3H), 1.15 (s, 3H), 1.00-1.02 (overlapping s and d, 6H)

EXAMPLE 56

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(4-(1-CARBOXYETHYL)PIPERIDINYLCARBONYL)-AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(4-(1-carboxyethyl)piperidinyldicarbonyl)amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (0.15 g; 0.23 mmol) in THF (5 mL) was added 1 M NaOH until a pH 10 solution persisted for 1 h. The

solution was evaporated under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

Anal: (C ₃₃ H ₅₂ N ₄ O ₆ S)					
Calc.	C, 53.09;	H, 6.65;	N, 6.84	1.6 TFA,	0.2 H ₂ O
Found	C, 53.08;	H, 6.66;	N, 6.85		

TLC: R_f 0.10 (80:20:2 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 8.72 min

FAB MS: m/z 633 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.38 (br s, 1H), 7.18 (m, 2H), 7.03 (m, 2H), 2.29 (s, 3H), 1.13 (s, 3H), 0.98-1.01 (overlapping s and d, 6H)

EXAMPLE 57

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(3-(1-ETHOXYCARBONYLMETHYL)-PIPERIDINYLCARBONYL)AMINO)PROPYL-BICYCLO(2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-*exo*-hydroxy-2-*endo*-2-(1-(3-piperidinylcarbonyl)amino)propyl-(2.2.1)bicycloheptan-1-yl)methanesulfonyl)-4-(2-methylphenyl)piperazine (0.20 g; 0.36 mmol) in DMF (5 mL) was added ethyl bromoacetate (0.044 mL; 0.40 mmol) and DIEA (0.070 mL; 0.40 mmol). After 24 h at ambient temperature, the solution was evaporated under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

Anal: (C ₃₄ H ₅₄ N ₄ O ₆ S)					
Calc.	C, 52.81;	H, 6.67;	N, 6.57	1.75 TFA,	0.35 H ₂ O
Found	C, 52.80;	H, 6.64;	N, 6.69		

TLC: R_f 0.35 (95:5 CHCl₃:MeOH)


HPLC (method A): retention time 9.26 min

FAB MS: m/z 647 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.19 (m, 2H), 7.04 (m, 2H), 4.26 (q, J = 7 Hz, 2H), 3.85 (s, 2H), 2.32 (s, 3H), 1.29 (t, J = 7 Hz, 3H), 1.14 (s, 3H), 1.02-1.05 (overlapping s and d, 6H)

EXAMPLE 58

1-((7,7-DIMETHYL-2-EXO-HYDROXY-2-ENDO-2-(1-(4-(1-CARBOXYMETHYL)PIPERIDINYLCARBONYL)-AMINO)PROPYL-BICYCLO (2.2.1)HEPTAN-1-YL)METHANESULFONYL)-4-(2-METHYLPHENYL)PIPERAZINE



To a stirred solution of 1-((7,7-dimethyl-2-exo-hydroxy-2-endo-2-(1-(3-(1-methoxycarbonyl)-piperidinylcarbonyl)amino)propyl-(2.2.1)bicycloheptan-1-yl) methanesulfonyl)-4-(2-methylphenyl)piperazine (0.15 g; 0.23 mmol) in THF (5 mL) was added 1 M NaOH until a pH 10 solution persisted for 1 h. The solution was evaporated under reduced pressure and the residue was purified by preparative reverse phase HPLC using an acetonitrile-water gradient containing 0.1% TFA. The TFA salt of title compound was obtained as a lyophilized powder.

Anal: (C ₃₂ H ₅₀ N ₄ O ₆ S)					
Calc.	C, 53.23;	H, 6.82;	N, 7.18	1.3 TFA,	0.75 H ₂ O
Found	C, 53.20;	H, 6.81;	N, 7.18		

TLC: R_f 0.15 (80:20:2 CHCl₃:MeOH:NH₄OH)


HPLC (method A): retention time 8.59 min

FAB MS: m/z 619 (M⁺ + H)

¹H NMR (400 MHz, CDCl₃): δ 7.35 (br s, 1H), 7.17 (m, 2H), 7.02 (m, 2H), 3.90 (s, 2H), 2.30 (s, 2H), 1.13 (s, 3H), 1.01 (s, 3H), 0.97 (d, J = 6 Hz, 3H)

TABLE

In addition to those compounds specifically exemplified above, additional compounds of the present invention are set forth in tabular form below. These compounds are synthesized by use of the synthetic routes and methods described in the above Schemes and Examples and variations thereof well known to those of ordinary skill in the art, and not requiring undue experimentation. All variables listed in the Tables below are with reference to the following generic structure:









TABLE


TABLE (Continued)

5




10

15




20

25




30

35



40

45




50

55


TABLE (Continued)

5




10

15




20

25




30

35



40

45




50

55


TABLE (Continued)

5



10


15



20


25

30



35

40



45


50

55

TABLE (Continued)


5

10




15

20




25

30




35



40

45



50

55

TABLE (Continued)

5

10

15

20

25

30

35

40

45

50

55






TABLE (Continued)

5




10

15




20

25




30



35

40



45


50

55

TABLE (Continued)

5


10



15


20

25




30

35



40

45




50


55

TABLE (Continued)


5




10




15




20




25



30



35









40

45

50

55

TABLE (Continued)

35

40

45

Additional examples of species covered by this invention include the following non-limiting list:

50

55

5

10

15

20

25

30


35

40


45

50

55




5



10


15



20

25

30



35

40


45

50

55

5


10



15

20


25



30

35

40



45 EXAMPLE 33

RADIOGLAND BINDING ASSAYS

50 The high affinity binding of [³H] Oxytocin (OT) ([tyrosyl, 3,5-[³H]OT; 30-60 Ci/mmol; New England Nuclear, Boston, MA) to uterine OT receptors was based on an assay using a crude membrane preparation of uteri taken from diethylstilbestrol dipropionate (DES)-treated (0.3 mg/kg, ip; 18-24) rats. Competition studies were conducted at equilibrium (60 minutes; 22° C) using 1 nM [³H]OT in the following assay buffer: 50 mM Tris-HCl, 5 mM MgCl₂, and 0.1% BSA, pH 7.4. Non-specific binding (10% of the total binding) was determined using 1 μM unlabeled OT and the binding reaction was terminated by filtration through glass fiber filters using a cell harvester (model 7019, Skatron, Inc., Sterling, VA). IC₅₀ (the concentration of tested compound that inhibits 50% of OT) was reported, unless otherwise noted.

* Fuchs, A-R; Fuchs, F; Soloff, MS. 1985 J. Clin. Endocrinol. Metab. 60:37.

The measurement of [³H]Vasopressin (AVP) ([phenylalanyl-3,4,5-³H]AVP; 80-90 Ci/mmol; N w England Nuclear) binding to a crude membrane preparation of male rat liver (AVP-V₁ sites) or kidney medulla (AVP-V₂ sites) was determined according to the method of Butlen, et al.^{**} Competition assays were conducted at equilibrium (30 minutes at 30°C) using 1 nM [³H]AVP (liver) or 2 nM [³H]AVP (kidney) in the following assay buffer: 100 mM Tris-HCl, 5 mM MgCl₂, 0.1% BSA, 50 μM phenylmethylsulfonylfluoride, and 50 μg/ml bacitracin, pH 8.0. Nonspecific binding (5-10% of the total binding) was determined using 10 μM unlabeled AVP, and the binding reaction was terminated by filtration as described above for the [³H]OT binding assay.

K_i values were obtained for each compound from three to six separate determinations of the IC₅₀ values ($K_i = IC_{50}/(1 + c/K_d)$)^{***} using K_d values obtained from saturation binding assay: [³H]OT (uterus), 0.7 nM; [³H]AVP (liver), 0.4 nM; [³H] (kidney), 1.4 nM.

	<u>Example</u>	<u>IC₅₀</u>
15	1	1,000 nM
	2	150 nM
	3	180 nM
20	4	34 nM
	5	100 nM
	6	10 nM
	7	8 nM
25	8	18 nM
	9	5 nM
	10	48% inhibition at 100 nM
30	11	54 nM
	12	23% inhibition at 100 nM

35

40

45

50

55 ** Butlen, D; Guillon, G; Rajerison, R.M.; Jard, S; Sawyer, W.H.; Manning, M. 1978 Mol Pharmacol 14:1006.

*** Cheng, Y-C; Prusoff, W.H.; 1973 Biochem Pharmacol 22:3099

ExampleIC₅₀

	14	1,100 nM
5	15	44% inhibition at 1,000 nM
	16	64% inhibition at 1,000 nM
	17	36% inhibition at 100 nM
	18	75% inhibition at 1,000 nM
10	19	31% inhibition at 1,000 nM
	20	72% inhibition at 1,000 nM
	21	38% inhibition at 1,000 nM
15	22	78% inhibition at 1,000 nM
	23	120 nM
	24	260 nM
20	25	34% inhibition at 100 nM
	26	35 nM
	27	37% inhibition at 100 nM
	28	35% inhibition at 100 nM
25	29	78% inhibition at 1,000 nM
	30	16% inhibition at 10,000 nM
	31	5% inhibition at 10,000 nM
30	32	37% inhibition at 1,000 nM
	33	460 nM
	34	-
35	35	91% inhibition at 100 nM
	36	7.7 nM
	37	1.2 nM
	38	5.4 nM
40	39	54% inhibition at 1,000 nM
	40	35% inhibition at 1,000 nM
	41	6.3 nM
45	42	9.2 nM
	43	110 nM

50


55

	<u>Example</u>	<u>IC₅₀</u>
	44	26 nM
5		180 nM
	45	12 nM
	46	20 nM
	47	15 nM
10	48	30 nM
	49	25 nM
	50	66% inhibition at 100 nM
15	51	38 nM
	52	66% inhibition at 100 nM
	53	28 nM
20	54	14 nM
	55	30 nM
	56	54 nM
25	57	66% inhibition at 100 nM
	58	56 nM

While the invention has been described and illustrated with reference to certain preferred embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the invention. For example, effective dosages other than the preferred dosages as set forth hereinabove may be applicable as a consequence of variations in the responsiveness of the mammal being treated for prevention of preterm labor, or for other indications for the compounds of the invention indicated above. Likewise, the specific pharmacological responses observed may vary according to and depending upon the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended, therefore, that the invention be limited only by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.

Claims

1. A compound of the formula:



and the pharmaceutically acceptable salts thereof, wherein

X is

- (1) carbonyl or
- (2) sulfonyl;

Y is

- (1) hydrogen,
- (2) alkyl or
- (3) NH;

Z is

- (1) C or
- (2) N;

R¹ and R² are independently one or more of

- (1) hydrogen,
- (2) halogen,
- (3) alkoxy,
- (4) alkylsulfonyl or
- (5) unsubstituted or substituted alkyl wherein said substituent is amino, alkyl, or dialkylamino;

R³ and R⁴ are independently one or more of

- (1) hydrogen,
- (2) alkyl,
- (3) substituted alkyl where said substituent is amino, alkylsulfonyl, arylsulfonyl, alkylamino or dialkylamino;
- (4) phenylalkyl or
- (5) oxo;

R⁵ is

- (1) hydrogen or
 (2) oxo;
 R⁶ and R⁷ are independently on or more of
 (1) hydrogen,
 5 (2) alkyl or
 (3) joined to form unsubstituted or substituted cycloalkyl where said substituent is
 hydroxy or
 hydroxyalkyl;
 R⁸, R⁹ and R¹⁰ are independently one or more of
 10 (1) hydrogen,
 (2) oxime,
 (3) hydroxy,
 (4) halogen,
 (5) cyclic epoxide,
 15 (6) alkylcarbonyloxy,
 (7) alkoxycarbonylalkoxy,
 (8) trihaloalkylsulfonyloxy,
 (9) unsubstituted or substituted 5 or 6-membered heterocyclic rings containing 1,2 or 3 heteroatoms
 where said heteroatom is N,S or O and said substituents are one or more of
 20 oxo or,
 C₁₋₆ alkyl, optionally substituted by
 alkylthio
 alkylsulfinyl
 alkylsulfonyl
 25 cyano
 carboxy,
 hydroxy
 amine,
 alkylamine
 30 dialkylamine
 alkoxy
 aminocarbonyl,
 tetrazolyl
 imidazole and
 35 spiro-piperidinyl where the piperidine
 N is unsubstituted or substituted by
 C₁₋₆ alkyl;
 (10) unsubstituted, mono- or di-substituted phenyl where said substituents are independently one or
 more of
 40 halogen,
 carboxy,
 alkoxycarbonyl,
 carboxyalkyl,
 amino,
 45 alkylamino,
 dialkylamino,
 cyano,
 alkylsulfonylamino,
 aminoalkyl,
 50 alkylaminoalkyl,
 dialkylaminoalkyl; and
 m, n, p and q are integers of from 0 to 2.

2. A compound of the formula:

55



30

35

40

45

50

55

77

(5) oxo;

R⁵ is

(1) hydrogen or

(2) oxo;

R⁶ and R⁷ are independently one or more of

(1) hydrogen,

(2) alkyl or

(3) joined to form unsubstituted or substituted cycloalkyl where said is substituent is hydroxy or

hydroxyalkyl;

R⁸, R⁹ and R¹⁰ are independently one or more of

(1) hydrogen,

(2) oxime,

(3) hydroxy,

(4) halogen,

(5) oxo,

(6) cyclic epoxide,

(7) alkylcarbonyloxy,

(8) alkoxy carbonylalkoxy,

(9) trihaloalkylsulfonyloxy,

(10) unsubstituted or substituted 5-membered heterocyclic rings containing 1 or 2 heteroatoms where said heteroatom is N or O and said substituents are oxo or

C₁₋₆ alkyl, optionally substituted by

carboxy,


amine,

aminocarbonyl or

imidazole; and

m, n, p and q are integers from 0 to 2.

3. A compound of the formula:



and the pharmaceutically acceptable salts thereof, wherein

R is

(1) hydrogen,

(2) oxime,

- (3) hydroxy,
 (4) halogen,
 (5) cyclic epoxide,
 (6) alkylcarbonyloxy,
 5 (7) alkoxycarbonylalkoxy,
 (8) trihaloalkylsulfonyloxy,
 (9) unsubstituted or substituted 5-membered heterocyclic rings containing 1 or 2 heteroatoms where said heteroatom is N or O and said substituents are independently one or more of
 oxo or,
 10 C₁₋₆ alkyl, optionally substituted by
 carboxy,
 amine,
 aminocarbonyl or,
 imidazole
- 15 4. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound as claimed in claim 1 in an amount sufficient to antagonize the binding of oxytocin to its receptor binding site.
- 20 5. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound as claimed in claim 1 in an amount sufficient to prevent preterm labor in a mammal.
6. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound as claimed in claim 1 in an amount sufficient to stop labor prior to cesarian delivery in a mammal.
- 25 7. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound as claimed in claim 1 in an amount sufficient to treat dysmenorrhea.
8. The use of a compound as claimed in claim 1 for the manufacture of a medicament for antagonizing the binding of oxytocin to its receptor binding site in a mammalian biologic system.
- 30 9. The use of a compound as claimed in claim 1 for the manufacture of a medicament for preventing preterm labor in a mammal.
- 35 10. The use of a compound as claimed in claim 1 for the manufacture of a medicament for stopping labor prior to cesarian delivery in a mammal.
- 40 11. The use of a compound as claimed in claim 1 for the manufacture of a medicament for treating dysmenorrhea in a mammal.

(19)



Eur päisches Patentamt
European Patent Office
Offic uropéen d s brevets



(11) Publication number:

0 533 242 A3

(12)

EUROPEAN PATENT APPLICATION(21) Application number: **92202722.2**(22) Date of filing: **08.09.92**

(51) Int. Cl.⁵: **C07D 295/22, C07D 207/26,
C07D 233/54, C07D 453/02,
C07D 303/34, C07D 211/60,
C07D 207/16, A61K 31/495**

(30) Priority: **16.09.91 US 760422**

(43) Date of publication of application:
24.03.93 Bulletin 93/12

(94) Designated Contracting States:
CH DE FR GB IT LI NL

(88) Date of deferred publication of the search report:
15.09.93 Bulletin 93/37


(71) Applicant: **MERCK & CO. INC.**
126, East Lincoln Avenue P.O. Box 2000
Rahway New Jersey 07065-0900(US)

(72) Inventor: **Bock, Mark G.**
1603 Leon Drive
Hatfield, PA 19440(US)
Inventor: **Williams, Peter D.**
260 Shady Nook Road
Harleysville, PA 19438(US)

(74) Representative: **Thompson, John Dr. et al**
Merck & Co., Inc. European Patent
Department Terlings Park Eastwick Road
Harlow, Essex CM20 2QR (GB)

(54) **Substituted derivatives of piperazinyllcamphorsulfonyl oxytocin antagonists.**

(57) Compounds of the formula:



ed oxo, where said substituent is unsubstituted or substituted saturated 6-membered heterocyclic rings having 1 or 2 heteroatoms which are N, and said substituent is alkyl.

The compounds of formula I are oxytocin antagonists useful in the treatment of preterm labor, dysmenorrhea and for the stoppage of labor preparatory to cesarean delivery. Also disclosed are pharmaceutical compositions containing these compounds, methods of their use and methods of their preparation.

wherein R is oxime, hydroxy, halogen, cyclic epoxide, alkylcarbonyloxy, alkoxy carbonylalkoxy, trihaloalkylsulfonyloxy, or unsubstituted or substitut-

EP 0 533 242 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 92 20 2722 - 2

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	US-A-4 147 870 (ANTONIO A. GARCIA) *Document*	1-3	C07D295/22 C07D207/26 C07D233/54 C07D453/02 C07D303/34 C07D211/60 C07D207/16 A61K31/495
A	--- CHEMICAL ABSTRACTS, vol. 86, no. 5, 31 January 1977, Columbus, Ohio, US; abstract no. 29877c, page 376 ;column R ; * abstract *	1-3	
A	& ES-A-418 775 (LABORATORIOS LIADE S. A.) ---		
A	FR-A-2 081 346 (LABORATORIOS LIADE S.A.) *Document*	1-3	
A	---		
A	FR-A-2 292 477 (LABORATORIOS LIADE S.A.) *Document*	1-3	
A	---		
A	CHEMICAL ABSTRACTS, vol. 78, no. 5, 5 February 1973, Columbus, Ohio, US; abstract no. 29818u, page 510 ;column L ; * abstract *	1-3	
	& ES-A-374 515 (LABORATORIOS LIADE) ---		
P,A	EP-A-0 486 280 (MERCK) *Document*	1-11	C07D
P,A	---		
P,A	EP-A-0 450 761 (MERCK) *Document*	1-11	

The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 12 JULY 1993	Examiner LUYTEN H.W.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure F : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document			